

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number  
WO 02/22592 A2

(51) International Patent Classification<sup>7</sup>: C07D 295/00

(74) Agent: LEE, William, Y.; Schering-Plough Corporation,  
Patent Department - K-6-1 1990, 2000 Galloping Hill Road,  
Kenilworth, NJ 07033-0530 (US).

(21) International Application Number: PCT/US01/28324

(22) International Filing Date:  
12 September 2001 (12.09.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,  
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,  
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT,  
RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ,  
VN, YU, ZA.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/232,255 14 September 2000 (14.09.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

(71) Applicant: SCHERING CORPORATION [US/US];  
Patent Department - K-6-1 1990, 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: GREENLEE, William, J.; 115 Herrick  
Avenue, Teaneck, NJ 07666 (US). HUANG, Ying; 8  
Ivy Terrace, East Brunswick, NJ 08816 (US). KELLY,  
Joseph, M.; 112 Princeton Road, Parlin, NJ 08859 (US).  
MC COMBIE, Stuart, W.; 28 Hanford Place, Caldwell,  
NJ 07006 (US). STAMFORD, Andrew, W.; 27 Overlook  
Road, Chatham Township, NJ 07928 (US). WU, Yusheng;  
235 East 40th Street, Apt. #20C, New York, NY 10016  
(US).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: SUBSTITUTED UREA NEUROPEPTIDE Y Y5 RECEPTOR ANTAGONISTS

(57) Abstract: Compounds represented by structural formula (I) including its N-oxides wherein Y is (I') R1 is H or (C1-C6)alkyl; R2 is H, (C1-C6)alkyl, (C3-C9)cycloalkyl or (C3-C7)cycloalkyl(C1-C6)alkyl; R3 is (II'); Z is OR10, -N(R9)R10 or -NH2; j is 0, 1 or 2; k is 1 or 2; l is 0, 1 or 2; m is 0, 1 or 2; R4 is 1-3 substituents independently selected from the group consisting of H, -OH, halogen, haloalkyl, (C1-C6)alkyl, (C3-C7)cycloalkyl, (C3-C7)cycloalkyl(C1-C6)alkyl, -CN, -O(C1-C6)alkyl, -O(C3-C7)cycloalkyl, -O(C1-C6)alkyl(C3-C7)cycloalkyl, -S(C1-C6)alkyl, -S(C3-C7)cycloalkyl, -S(C1-C6)alkyl(C3-C7)cycloalkyl, -NH2, -NR9R10, -NO2, -CONH2, -CONR9R10 and NR2COR10; R5 is 1-3 substituents independently selected from the group consisting of H, halogen, -OH, haloalkyl, haloalkoxy, -CN, -NO2, (C1-C6)alkyl, (C3-C7)cycloalkyl, (C3-C7)cycloalkyl(C1-C6)alkyl, -O(C1-C6)alkyl, -O(C3-C7)cycloalkyl, -O(C1-C6)alkyl(C3-C7)cycloalkyl, -CONH2 and -CONR9R10; R6 is -SO2(C1-C6)alkyl, -SO2(C3-C7)cycloalkyl, -SO2(C1-C6)alkyl(C3-C7)cycloalkyl, -SO2(C1-C6)haloalkyl, -SO2(hydroxy(C2-C6)alkyl), -SO2(amino(C2-C6)alkyl), -SO2(alkoxy(C2-C6)alkyl), -SO2(alkylamino(C2-C6)alkyl), -SO2(di-alkylamino(C2-C6)alkyl), -SO2(aryl), -SO2(heteroaryl), -SO2(aryl(C2-C6-alkyl)), SO2NH2, -SO2NR9R10, -C(O)C1-C6alkyl, -C(O)C3-C7cycloalkyl, -C(O)aryl, -C(O)heteroaryl, -C(O)NR9R10, -C(O)NH2, -C(S)NR9R10, -C(S)NH2, aryl, heteroaryl, -(CH2)nC(O)NH2, -(CH2)nC(O)NR9R10, -C(=NCN)alkylthio, -C(=NCN)NR9R10, (C1-C6)alkyl, (C3-C7)cycloalkyl, (C3-C7)cycloalkyl(C1-C6)alkyl, aryl(C1-C6)alkyl, heteroaryl(C1-C6)alkyl or -C(O)OR9, n= 1 to 6; R7 = H or alkyl; R8 is H, (C1-C6)alkyl, (C3-C7)cycloalkyl, (C3-C7)cycloalkyl(C1-C6)alkyl, aryl, heteroaryl, -SO2(C1-C6)alkyl, -SO2(C3-C7)cycloalkyl, -SO2(C1-C6)alkyl(C3-C7)cycloalkyl, -SO2(C1-C6)haloalkyl or -SO2(aryl); R9 is (C1-C6)alkyl, (C3-C7)cycloalkyl, (C3-C7)cycloalkyl(C1-C6)alkyl, aryl(C1-C6)alkyl, aryl or heteroaryl; and, R10 is hydrogen, (C1-C6)alkyl, (C3-C7)cycloalkyl, (C3-C7)cycloalkyl(C1-C6)alkyl, aryl(C1-C6)alkyl, aryl or heteroaryl; or a pharmaceutically acceptable addition salt and/or hydrate thereof, or prodrug thereof, or R9 and R10 taken together can form a 4-7 membered ring containing 1 or 2 heteroatoms; or where applicable, a geometric or optical isomer or a racemic mixture thereof, are claimed, as well as additional novel compounds; also claimed are pharmaceutical compositions and methods of using the aforesaid compounds in the treatment of obesity, eating disorders such as hyperphagia and diabetes.

WO 02/22592 A2

SUBSTITUTED UREA  
NEUROPEPTIDE Y Y5 RECEPTOR ANTAGONISTS

5

Field of the Invention

The present invention relates to neuropeptide Y Y5 receptor antagonists useful in the treatment of obesity and eating disorders, pharmaceutical compositions containing the compounds, and methods of treatment using the compounds.

10

Background of the Invention

Neuropeptide Y (NPY) is a 36 amino acid neuropeptide that is widely distributed in the central and peripheral nervous systems. NPY is a member of the pancreatic polypeptide family that also includes peptide YY and pancreatic polypeptide (Wahlestedt, C., and Reis, D., Ann. Rev. Toxicol., 32, 309, 1993). NPY elicits its physiological effects by activation of at least six receptor subtypes designated Y1, Y2, Y3, Y4, Y5 and Y6 (Gehlert, D., Proc. Soc. Exp. Biol. Med., 218, 7, 1998; Michel, M. et al., Pharmacol. Rev., 50, 143, 1998). Central administration of NPY to animals causes dramatically increased food intake and decreased energy expenditure (Stanley, B. and Leibowitz, S., Proc. Natl. Acad. Sci. USA 82: 3940, 1985; Billington et al., Am J. Physiol., 260, R321, 1991). These effects are believed to be mediated at least in part by activation of the NPY Y5 receptor subtype. The isolation and characterization of the NPY Y5 receptor subtype has been reported (Gerald, C. et al., Nature, 1996, 382, 168; Gerald, C. et al. WO 96/16542). Additionally, it has been reported that activation of the NPY Y5 receptor by administration of the Y5 – selective agonist [D-Trp<sup>32</sup>]NPY to rats stimulates feeding and decreases energy expenditure (Gerald, C. et al., Nature, 1996, 382, 168; Hwa, J. et al., Am. J. Physiol., 277 (46), R1428, 1999). Hence, compounds that block binding of NPY to the NPY Y5 receptor subtype should have utility in the treatment of obesity, disorders such as, bulimia nervosa, anorexia nervosa, and in the treatment of disorders associated with obesity such as type II diabetes, insulin resistance, hyperlipidemia, and hypertension.

Published PCT patent application WO 00/27845 describes a class of compounds, characterized therein as spiro-indolines, said to be selective neuropeptide Y Y5 receptor antagonists and useful for the treatment of obesity and the complications associated therewith. Known urea derivatives indicated as


35

possessing therapeutic activity are described in U.S. Patent Nos. 4,623,662 (antiatherosclerotic agents) and 4,405,644 (treatment of lipometabolism). Provisional application, U.S. Serial No. 60/232,255 describes a class of substituted urea neuropeptide Y Y5 receptor antagonists.

5


### SUMMARY OF THE INVENTION

The present invention relates to compounds represented by the structural formula I:




10 including its N-oxides, wherein

Y is




R<sup>1</sup> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sup>2</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>9</sub>)cycloalkyl or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;



15



Z is OR<sup>10</sup>, -N(R<sup>9</sup>)(R<sup>10</sup>) or -NH<sub>2</sub>;

j is 0, 1 or 2;

20

k is 1 or 2;

l is 0, 1 or 2;

m is 0, 1 or 2;

R<sup>4</sup> is 1-3 substituents independently selected from the group consisting of H, -OH, halogen, haloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,

-CN, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
 -S(C<sub>1</sub>-C<sub>6</sub>)alkyl, -S(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -S(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -NH<sub>2</sub>, -NR<sup>9</sup>R<sup>10</sup>,  
 -NO<sub>2</sub>, -CONH<sub>2</sub>, -CONR<sup>9</sup>R<sup>10</sup> and NR<sup>2</sup>COR<sup>10</sup>;

R<sup>5</sup> is 1-3 substituents independently selected from the group consisting of H,  
 5 halogen, -OH, haloalkyl, haloalkoxy, -CN, -NO<sub>2</sub>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
 (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
 -O(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -CONH<sub>2</sub> and -CONR<sup>9</sup>R<sup>10</sup>;

R<sup>6</sup> is -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO<sub>2</sub>(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
 -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)haloalkyl, -SO<sub>2</sub>(hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>(amino(C<sub>2</sub>-C<sub>6</sub>)alkyl),  
 10 -SO<sub>2</sub>(alkoxy(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>(alkylamino(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>(dialkylamino(C<sub>2</sub>-C<sub>6</sub>)alkyl),  
 -SO<sub>2</sub>(aryl), -SO<sub>2</sub>(heteroaryl), -SO<sub>2</sub>(aryl(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>,  
 -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -C(O)(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 -C(O)aryl, -C(O)heteroaryl, -C(O)NR<sup>9</sup>R<sup>10</sup>, -C(O)NH<sub>2</sub>, -C(S)NR<sup>9</sup>R<sup>10</sup>, -C(S)NH<sub>2</sub>, aryl,  
 heteroaryl, -(CH<sub>2</sub>)<sub>n</sub>C(O)NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>n</sub>C(O)NR<sup>9</sup>R<sup>10</sup>, -C(=NCN)alkylthio, -  
 15 C(=NCN)NR<sup>9</sup>R<sup>10</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-  
 C<sub>6</sub>)alkyl, heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or -C(O)OR<sup>9</sup>, n= 1 to 6;

R<sup>7</sup> = H or alkyl;

R<sup>8</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl,  
 heteroaryl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO<sub>2</sub>(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
 20 -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)haloalkyl or -SO<sub>2</sub>(aryl);

R<sup>9</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or heteroaryl; and,

R<sup>10</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
 aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or heteroaryl;  
 25 or R<sup>9</sup> and R<sup>10</sup> taken together can form a 4-7 membered ring containing 1 or 2  
 heteroatoms;

or a pharmaceutically acceptable addition salt and/or hydrate thereof, or  
 prodrug thereof, or where applicable; a geometric or optical isomer or a racemic  
 mixture thereof.

30 The present invention also relates to a method of treating obesity and eating  
 disorders, such as hyperphagia, and diabetes comprising administering to a mammal  
 in need of such treatment an effective amount of a compound of formula I.

Another aspect of the invention is a pharmaceutical composition for treating obesity, eating disorders and diabetes which comprises a compound of formula I in combination with a pharmaceutically acceptable carrier.

## 5 DETAILED DESCRIPTION

Except where stated otherwise, the following definitions apply throughout the present specification and claims. These definitions apply regardless of whether a term is used by itself or in combination with other terms. Hence the definition of "alkyl" applies to "alkyl" as well as to the "alkyl" portions of "alkoxy", etc.

10 Alkyl represents a straight or branched saturated hydrocarbon chain having the designated number of carbon atoms. Where the number of carbon atoms is not specified, 1 to 6 carbons are intended.

Halo represents fluoro, chloro, bromo or iodo.

Haloalkyl refers to alkyl substituted by halo, wherein the number of halo  
15 substituents ranges from one to as many halo substituents required for full substitution of the alkyl substituent.

Aryl refers to a mono- or bicyclic ring system having at least one aromatic ring including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, and the like. The aryl group can be unsubstituted or substituted with one, two, or three substituents  
20 independently selected from lower alkyl, halo, cyano, nitro, haloalkyl, hydroxy, alkoxy, carboxy, carboxamide, mercapto, sulfhydryl, amino, alkylamino and dialkylamino.

Heteroaryl refers to 5- to 10-membered single or benzofused aromatic rings consisting of 1 to 3 heteroatoms independently selected from the group consisting of -O-, -S-, and -N=, provided that the rings do not possess adjacent oxygen and sulfur  
25 atoms. The heteroaryl group can be unsubstituted or substituted with one, two, or three substituents independently selected from lower alkyl, halo, cyano, nitro, haloalkyl, hydroxy, alkoxy, carboxy, carboxamide, mercapto, sulfhydryl, amino, alkylamino, dialkylamino.

When a variable appears more than once in the structural formula, for example  
30 R<sup>9</sup>, the identity to each variable appearing more than once may be independently selected from the definition for that variable.


N-oxides can form on a tertiary nitrogen present in an R substituent, or on =N- in a heteroaryl ring substituent and are included in the compounds of formula I.

For compounds of the invention having at least one asymmetrical carbon atom, all isomers, including diastereomers, enantiomers and rotational isomers are contemplated as being part of this invention. The invention includes d and l isomers in both pure form and in admixture, including racemic mixtures. Isomers can be prepared using conventional techniques, either by separating isomers of a compound of formula I or by synthesizing individual isomers of a compound of formula I.

Compounds of formula I can exist in unsolvated and solvated forms, including hydrated forms. In general, the solvated forms, with pharmaceutically acceptable solvents such as water, ethanol and the like, are equivalent to the unsolvated forms for purposes of this invention.


A compound of formula I may form pharmaceutically acceptable salts with organic and inorganic acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. The salts are prepared by contacting the free base forms with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution, such as dilute aqueous sodium hydroxide, potassium carbonate, ammonia or sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the salts are otherwise equivalent to their respective free base forms for purposes of the invention.

In a preferred group of compounds of formula 1, Y is



including, in particular, those compounds in which  $R^5$  is 1-3 substituents independently selected from the group consisting of H, halogen, haloalkyl and haloalkoxy and the sum of j and k is 1, 2 or 3.

In another preferred group of compounds of formula 1, Y is




- including, in particular, those compounds in which  $R^5$  and  $R^6$  each independently is 1 to 3 substituents independently selected from the group consisting of H, halogen, haloalkyl and haloalkoxy and the sum of j and k is 1, 2 or 3.

Compounds of formula I may be produced by processes known to those skilled in the art as shown in the following reaction schemes and in the preparations and examples below.

7

Scheme 1




- In Scheme 1, a 4-halophenyl isocyanate is condensed with an amino substituted cyclic amine derivative to give a 4-halophenyl urea derivative. Cleavage of the cyclic amine protecting group by methods known to those skilled in the art affords a cyclic amine derivative that can be derivatized, for example by alkylation (Path 1). Coupling of the product with, for example, an arylboronic acid, under palladium catalysis (Suzuki coupling) yields a biaryl urea derivative. Alternatively, the condensation product can be arylated, for example, by use of a Suzuki coupling



reaction (Path 2). When A is a protecting group, deprotection affords an amine that can be derivatized by, for example, sulfonylation, acylation or alkylation.


Scheme 2



In Scheme 2, reaction of an aryl lithium, for example, 5-thienyl lithium, with trimethylborate and coupling of the resultant boronate with a 4-haloaniline under palladium catalysis yields a biaryl amine derivative. Protection of the amine with, for example, trifluoroacetic anhydride gives a trifluoroacetamide derivative that can be halogenated with an appropriate halogenating agent, for example  $N$ -chlorosuccinimide. The protecting group can be cleaved and the resultant amine can be reacted with, for example,  $N,N'$ -disuccinimidyl carbonate and an amino substituted cyclic amine derivative, for example an amino piperidine derivative, to give a substituted urea. Cleavage of the piperidine nitrogen protecting group gives an amine that can be derivatized, for example, by sulfonylation or acylation.

9

Scheme 3



5

In Scheme 3, a 4-haloaniline or 4-halonitrobenzene derivative is arylated by use of, for example, a Suzuki coupling reaction. When X is a nitro group, the nitro group is subsequently reduced to an amine. The biaryl amine derivative can be

converted to an isocyanate derivative, which can be condensed with an amino substituted cyclic amine derivative (Path 3). Alternatively, condensation with an amino substituted cycloalkyl derivative affords cycloalkyl urea derivatives (Paths 4 and 5). An appropriately functionalized cycloalkyl urea derivative can be further functionalized as shown, for example, in Path 5.

The compounds of formula I exhibit selective neuropeptide Y Y5 receptor antagonizing activity, which has been correlated with pharmaceutical activity for treating obesity, eating disorders, such as hyperphagia, and diabetes.

Another aspect of this invention is a method of treating a mammal (e.g., human) having a disease or condition mediated by the neuropeptide Y Y5 receptor by administering a therapeutically effective amount of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug to the mammal.

Another aspect of this invention is directed to a method of treating obesity comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of Formula I or a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug.

Another aspect of this invention is directed to a method for treating metabolic and eating disorders such as bulimia and anorexia comprising administering to a mammal a therapeutically effective amount of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug.

Another aspect of this invention is directed to a method for treating hyperlipidemia comprising administering to a mammal a therapeutically effective amount of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug.

Another aspect of this invention is directed to a method for treating cellulite and fat accumulation comprising administering to a mammal a therapeutically effective amount of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug.

Another aspect of this invention is directed to a method for treating Type II diabetes comprising administering to a mammal a therapeutically effective amount of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug.

In addition to the "direct" effect of the compounds of this invention on the neuropeptide Y Y5 receptor subtype, there are diseases and conditions that will benefit from the weight loss such as insulin resistance, impaired glucose tolerance, Type II Diabetes, hypertension, hyperlipidemia, cardiovascular disease, gall stones, certain cancers, and sleep apnea.

This invention is also directed to pharmaceutical compositions, which comprise an amount of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug and a pharmaceutically acceptable carrier therefor.

This invention is also directed to pharmaceutical compositions for the treatment of obesity which comprise an obesity treating amount of a compound of Formula, I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug and a pharmaceutically acceptable carrier therefor.

Compounds of Formula I can be produced by processes known to those skilled in the art using either solution phase or solid phase synthesis as shown in the following reaction schemes, in the preparations and examples below.

The compounds of formula I display pharmacological activity in test procedures designed to demonstrate neuropeptide Y Y5 receptor antagonist activity. The compounds are non-toxic at pharmaceutically therapeutic doses. Following are descriptions of the test procedures.

#### cAMP Assay

HEK-293 cells expressing the Y5 receptor subtype were maintained in Dulbecco's modified Eagles' media (Gico-BRL) supplemented with 10% FCS (ICN), 1% penicillin-streptomycin and 200 µg/ml Geneticin® (GibcoBRL #11811-031) under a humidified 5% CO<sub>2</sub> atmosphere. Two days prior to assay, cells were released from T-175 tissue culture flasks using cell dissociation solution (1X; non-enzymatic [Sigma #C-5914]) and seeded into 96-well, flat-bottom tissue culture plates at a density of 15,000 to 20,000 cells per well. After approximately 48 hours, the cell monolayers were rinsed with Hank's balanced salt solution (HBSS) then preincubated with approximately 150 µl/well of assay buffer (HBSS supplemented with 4 mM MgCl<sub>2</sub>, 10 mM HEPES, 0.2% BSA [HH]) containing 1 mM 3-isobutyl-1-methylxanthine ([IBMX] Sigma #1-587) with or without the antagonist compound of interest at 37°C. After

20 minutes the 1 mM IBMX-HH assay buffer ( $\pm$  antagonist compound) was removed and replaced with assay buffer containing 1.5  $\mu$ M (CHO cells) or 5  $\mu$ M (HEK-293 cells) forskolin (Sigma #F-6886) and various concentrations of NPY in the presence or absence of one concentration of the antagonist compound of interest. At the end of 10 minutes, the media were removed and the cell monolayers treated with 75  $\mu$ l ethanol. The tissue culture plates were agitated on a platform shaker for 15 minutes, after which the plates were transferred to a warm bath in order to evaporate the ethanol. Upon bringing all wells to dryness, the cell residues were resolubilized with 250  $\mu$ l FlashPlate® assay buffer. The amount of cAMP in each well was quantified using the [ $^{125}$ I]-cAMP FlashPlate® kit (NEN #SMP-001) and according to the protocol provided by the manufacturer. Data were expressed as either pmol cAMP/ml or as percent of control. All data points were determined in triplicate and EC<sub>50</sub>'s (nM) were calculated using a nonlinear (sigmoidal) regression equation (GraphPad Prism™). The K<sub>B</sub> of the antagonist compound was estimated using the following formula:

$$K_B = [B] / (1 - \{[A'] / [A]\})$$

where [A] is the EC<sub>50</sub> of the agonist (NPY) in the absence of antagonist,

[A'] is the EC<sub>50</sub> of the agonist (NPY) in the presence of antagonist,

and [B] is the concentration of the antagonist.

#### NPY Receptor Binding Assay

Human NPY Y5 receptors were expressed in CHO cells. Binding assays were performed in 50 mM HEPES, pH 7.2, 2.5 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub> and 0.1% BSA containing 5-10  $\mu$ g of membrane protein and 0.1 nM  $^{125}$ I-peptide YY in a total volume of 200  $\mu$ l. Non-specific binding was determined in the presence of 1  $\mu$ M NPY. The reaction mixtures were incubated for 90 minutes at room temperature then filtered through Millipore MAFC glass fiber filter plates which had been pre-soaked in 0.5% polyethyleneimine. The filters were washed with phosphate-buffered saline, and radioactivity was measured in a Packard TopCount scintillation counter.

For the compounds of this invention, a range of neuropeptide Y5 receptor binding activity from about 0.2 nM to about 500 nM was observed. Compounds of this invention preferably have a binding activity in the range of about 0.2 nM to 250 nM, more preferably about 0.2 to 100 nM, and most preferably about 0.2 to 10 nM.

Yet another aspect of this invention are combinations of a compound of Formula I, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug and other compounds as described below.

Accordingly, another aspect of this invention is a method for treating obesity comprising administering to a mammal (e.g., a female or male human)

5       a.     an amount of a first compound, said first compound being a Formula I compound, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug; and

10       b.     an amount of a second compound, said second compound being an anti-obesity and/or anorectic agent such as a  $\beta_3$  agonist, a thyromimetic agent, an anorectic agent, or an NPY antagonist wherein the amounts of the first and second compounds result in a therapeutic effect.

This invention is also directed to a pharmaceutical combination composition comprising: a therapeutically effective amount of a composition comprising

15       a first compound, said first compound being a Formula I compound, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug

      a second compound, said second compound being an anti-obesity and/or anorectic agent such as a  $\beta_3$  agonist, a thyromimetic agent, an anorectic, or an NPY antagonist; and/or optionally a pharmaceutical carrier, vehicle or diluent.

20       Another aspect of this invention is a kit comprising:

      a.     an amount of a Formula I compound, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug and a pharmaceutically acceptable carrier, vehicle or diluent in a first unit dosage form;

      b.     an amount of an anti-obesity and/or anorectic agent such as a  $\beta_3$

25       agonist, a thyromimetic agent, an anorectic agent, or an NPY antagonist and a pharmaceutically acceptable carrier, vehicle or diluent in a second unit dosage form; and

      c.     means for containing said first and second dosage forms wherein the amounts of the first and second compounds result in a therapeutic effect.

30       Preferred anti-obesity and/or anorectic agents (taken singly or in any combination thereof) in the above combination methods, combination compositions and combination kits are:

phenylpropanolamine, ephedrine, pseudoephedrine, phentermine, a cholecystokinin-A (hereinafter referred to as CCK-A) agonist, a monoamine reuptake inhibitor (such as sibutramine), a sympathomimetic agent, a serotonergic agent (such as dexfenfluramine or fenfluramine), a dopamine agonist (such as bromocriptine), a melanocyte-stimulating hormone receptor agonist or mimetic, a melanocyte-stimulating hormone analog, a cannabinoid receptor antagonist, a melanin concentrating hormone antagonist, the OB protein (hereinafter referred to as "leptin"), a leptin analog, a leptin receptor agonist, a galanin antagonist or a GI lipase inhibitor or decreaser (such as orlistat). Other anorectic agents include bombesin agonists, dehydroepiandrosterone or analogs thereof, glucocorticoid receptor agonists and antagonists, orexin receptor antagonists, urocortin binding protein antagonists, agonists of the glucagon-like peptide-1 receptor such as Exendin and ciliary neurotrophic factors such as Axokine.

Another aspect of this invention is a method treating diabetes comprising administering to a mammal (e.g., a female or male human)

- a. an amount of a first compound, said first compound being a Formula I compound, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug; and
- b. an amount of a second compound, said second compound being an aldose reductase inhibitor, a glycogen phosphorylase inhibitor, a sorbitol dehydrogenase inhibitor, a protein tyrosine phosphatase 1B inhibitor, a dipeptidyl protease inhibitor, insulin (including orally bioavailable insulin preparations), an insulin mimetic, metformin, acarbose, a PPAR-gamma ligand such as troglitazone, rosiglitazone, pioglitazone or GW-1929, a sulfonylurea, glipazide, glyburide, or chlorpropamide wherein the amounts of the first and second compounds result in a therapeutic effect.

This invention is also directed to a pharmaceutical combination composition comprising: a therapeutically effective amount of a composition comprising a first compound, said first compound being a Formula I compound, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug; a second compound, said second compound being an aldose reductase inhibitor, a glycogen phosphorylase inhibitor, a sorbitol dehydrogenase inhibitor, a protein tyrosine phosphatase 1B inhibitor, a dipeptidyl protease inhibitor, insulin

(including orally bioavailable insulin preparations), an insulin mimetic, metformin, acarbose, a PPAR-gamma ligand such as troglitazone, rosiglitazone, pioglitazone, or GW-1929, a sulfonylurea, glipazide, glyburide, or chlorpropamide; and optionally a pharmaceutical carrier, vehicle or diluent.

5 Another aspect of this invention is a kit comprising:

- a. an amount of a Formula I compound, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug and a pharmaceutically acceptable carrier, vehicle or diluent in a first unit dosage form;
- b. an amount of an aldose reductase inhibitor, a glycogen phosphorylase  
10 inhibitor, a sorbitol dehydrogenase inhibitor, a protein tyrosine phosphatase 1B inhibitor, a dipeptidyl protease inhibitor, insulin (including orally bioavailable insulin preparations), an insulin mimetic, metformin, acarbose, a PPAR-gamma ligand such as troglitazone, rosiglitazone, pioglitazone, or GW-1929, a sulfonylurea, glipazide, glyburide, or chlorpropamide and a pharmaceutically acceptable carrier, vehicle or  
15 diluent in a second unit dosage form; and
- c. means for containing said first and second dosage forms wherein the amounts of the first and second compounds result in a therapeutic effect.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid.

- 20 Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g., magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral  
25 administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 18<sup>th</sup> Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

- 30 Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.



Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations which are intended to be converted,  
5 shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal composition can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type  
10 as are conventional in the art for this purpose.

Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate quantities of the active component, e.g., an effective amount to achieve  
15 the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.01 mg to about 1000 mg, preferably from about 0.01 mg to about 750 mg, more preferably from about 0.01 mg to about 500 mg, and most preferably from about 0.01 mg to about 250 mg, according to the particular  
20 application.

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total dosage may be divided and administered in portions during the  
25 day as required.


The amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical  
30 recommended daily dosage regimen for oral administration can range from about 0.04 mg/day to about 4000 mg/day, in two to four divided doses.

The invention disclosed herein is exemplified by the following preparations and examples which should not be construed to limit the scope of the disclosure.

Alternative mechanistic pathways and analogous structures may be apparent to those skilled in the art.

In the preparations and examples, the following abbreviations are used: room temperature (R.T.), phenyl(Ph), -t-butyloxycarbonyl(-Boc), methylamine (MeNH<sub>2</sub>), sodium triacetoxyborohydride (NaBH(OAc)<sub>3</sub>), ethyl acetate (EtOAc), methanol (MeOH), triethylamine (Et<sub>3</sub>N), ether (Et<sub>2</sub>O), tetrahydrofuran (THF), diisopropylethylamine (iPr<sub>2</sub>NEt), 1,2-dimethoxyethane (DME), ethanol (EtOH) and preparative thin layer chromatography (PTLC).



10

Preparation 1

15


To a mixture of N-t-butyloxycarbonyl-4-piperidone (10.0 g, 50 mmol) and aqueous methylamine (40% w/w, 10 ml) in 1,2-dichloroethane (125 ml) was added NaBH(OAc)<sub>3</sub> (16.0 g, 75 mmol). The reaction mixture was stirred overnight, then 1M NaOH (250 ml) was added and the whole was extracted with ether (700 ml). The organic layer was washed with sat'd NaCl, dried (MgSO<sub>4</sub>), filtered, and concentrated to give the product (10.5 g, 97%) as an oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 4.09 (2H, m), 2.86 (2H, m), 2.55 (1H, m), 2.50 (3H, s), 1.90 (2H, m), 1.51 (9H, s), 1.30 (2H, m).

20

Preparation 2Step 1

To a mixture of N-benzyloxycarbonyl-4-piperidone (10.70 g, 43.1 mmol) and aq. 40% MeNH<sub>2</sub> (6.67 g, 85.8 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (200 ml) at R.T. was added NaBH(OAc)<sub>3</sub> (27.25 g, 128.6 mmol). The reaction mixture was stirred at R.T. for 3 h then poured into sat'd NaHCO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x200 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to give the product (10.63 g, 100%) that was used without further purification. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.34 (5H, m), 5.12 (2H, s), 4.19 (2H, b), 2.87 (2H, b), 2.72 (1H, m), 2.49 (3H, s), 1.92 (2H, b), 1.42 (2H, m). MS m/e 249 (M+H).

### 10 Step 2




To the product of Step 1 (10.63 g, 42.9 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (200 ml) at R.T. was added di-*tert*-butyl dicarbonate (11.30 g, 51.8 mmol) in portions. The reaction mixture was allowed to stir at R.T. for 5 h then poured into 1N NaOH (50 ml)/CH<sub>3</sub>OH (10 ml). The mixture was stirred for 15 min. and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x200 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The residue was subjected to column chromatography (gradient 1:10 to 1:4 EtOAc/hexane) to give the product (13.00 g, 87%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.33 (5H, m), 5.10 (2H, s), 4.19 (3H, m), 2.87 (2H, b), 2.68 (3H, s), 1.60 (4H, m), 1.44 (9H, s). MS m/e 349 (M+H).


### Step 3

A mixture of the product of Step 2 (12.90 g, 37.0 mmol) and 10% Pd/C (1.29 g) in MeOH (300 ml) was stirred under an H<sub>2</sub> atmosphere. After 16 h the reaction mixture was filtered through celite and the filter pad was washed with MeOH. The combined filtrate and washings were concentrated to afford the product (7.80 g, 98.3%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 4.19 (1H, b), 3.15 (2H, b), 2.74 (3H, s), 2.66 (2H, m), 1.63 (4H, m), 1.46 (9H, s). MS m/e 215 (M+H).

19


Preparation 3

To a stirred solution of Preparation 1 (21.0 g, 83.7 mmol) and Et<sub>3</sub>N (35 ml, 252 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (300 ml) was added benzyl chloroformate (18 ml, 126 mmol) dropwise. After 5 h, sat'd NH<sub>4</sub>Cl (200 ml) was added, and the organic layer was washed with H<sub>2</sub>O (150 ml) and sat'd NaCl (150 ml), dried (MgSO<sub>4</sub>), filtered and concentrated. To the residue (32 g) was added 4N HCl in 1,4-dioxane (300 ml), and the mixture was stirred for 4 h. The reaction mixture was concentrated, acetone was added, and the reaction mixture was again concentrated. The solid residue was dissolved in MeOH (40 ml) and Et<sub>2</sub>O was added. The resultant precipitate was collected, washed with Et<sub>2</sub>O, and dried to give the product as a white solid (20.2 g, 85%). MS m/e 249 (M+H, free base).

Example 1

15

1

Step 1


1-1-1

20

To a solution of Preparation 1 (7.0 g, 33 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (200 ml) was added 4-bromophenyl isocyanate (6.8 g, 35 mmol). The reaction mixture was stirred for 16 h, then H<sub>2</sub>O (200 ml) was added, and the organic layer was dried (MgSO<sub>4</sub>), filtered and evaporated. The residue was triturated with hexanes to give a white solid (11.0 g, 81%). MS (FAB) m/e 411 (M+H)<sup>+</sup>.


25

20


Step 2

To a solution of the product of Step 1 (400 mg, 0.97 mmol) and Pd(dppf)Cl<sub>2</sub>·CH<sub>2</sub>Cl<sub>2</sub> (200 mg, 0.24 mmol) in toluene (10 ml) was added 2-fluorophenylboronic acid (250 mg, 1.43 mmol), Cs<sub>2</sub>CO<sub>3</sub> (350 mg, 1.1 mmol), and H<sub>2</sub>O (0.3 ml). The reaction mixture was heated in a 90 °C oil bath under N<sub>2</sub> for 1 h, then allowed to cool. The reaction mixture was partitioned between EtOAc (100 ml) and H<sub>2</sub>O (50 ml). The organic layer was dried (MgSO<sub>4</sub>), filtered and evaporated. Flash chromatography (3:7 acetone/hexane) of the residue afforded the product (400 mg, 97%). HRMS calc. for C<sub>24</sub>H<sub>31</sub>FN<sub>3</sub>O<sub>3</sub> (M+H) 428.2349. Found 428.2343.

Coupling of the product of Step 1 with the appropriate boronic acid by essentially the same procedure gave:





HRMS calc. for C<sub>25</sub>H<sub>31</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub> (M+H) 478.2318. Found 478.2313.





HRMS calc. for C<sub>25</sub>H<sub>31</sub>F<sub>3</sub>N<sub>3</sub>O<sub>3</sub> (M+H) 478.2318. Found 478.2313.

20


21

HRMS calc. for  $C_{25}H_{31}F_3N_3O_4$  (M+H) 494.2260. Found 494.2267.

5

HRMS calc. for  $C_{24}H_{31}FN_3O_3$  (M+H) 428.2343. Found 428.2349.MS (FAB) m/e 478 (M+H)<sup>+</sup>.


10

MS (FAB) m/e 446 (M+H)<sup>+</sup>.Step 3

To a solution of the product of Step 2 (100 mg, 0.23 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 ml) was added 4 M HCl in 1,4-dioxane (3 ml). After 16 h, the reaction mixture was concentrated. The residue was triturated with ether and the solid was collected, washed with ether, and air-dried to give the product (80 mg, 96%). HRMS calc. for


5  $\text{C}_{19}\text{H}_{23}\text{FN}_3\text{O}$  (M+H) 328.1825. Found 328.1823.

Treatment of the other products from Step 2 by essentially the same procedure gave:




1-3-2

10 MS (ES) m/e 378 (M+H)<sup>+</sup>.



1-3-3


MS (FAB) m/e 378 (M+H)<sup>+</sup>.



1-3-4

15


HRMS calc. for  $\text{C}_{20}\text{H}_{23}\text{F}_3\text{N}_3\text{O}_2$  (M+H) 394.1742. Found 394.1747.



1-3-5

HRMS calc. for  $\text{C}_{19}\text{H}_{23}\text{FN}_3\text{O}$  (M+H) 328.1825. Found 328.1823.


20



1-3-6

MS (ES) m/e 378 (M+H)<sup>+</sup>.

23




1-3-7

HRMS calc. for  $C_{19}H_{22}F_2N_3O$  (M+H) 346.1731. Found 346.1725.




#### Step 4



- 5 To a stirred solution of the product of Step 3 (20 mg, 0.055 mmol) and triethylamine (0.1 ml, 0.7 mmol) in  $CH_2Cl_2$  (10 ml) was added methanesulfonyl chloride (0.1 ml, 0.1 mmol). After 16 h the reaction mixture was concentrated and the residue was subjected to PTLC (1:2 acetone/hexanes) to give a white solid (15 mg, 67%). HRMS calc. for  $C_{20}H_{25}FN_3O_3S$  (M+H) 406.1601. Found 406.1599.
- 10 The following examples were prepared from the appropriate starting amine and sulfonyl chloride.



Y	R <sup>6</sup>	MS (M+H)	Example
	-SO <sub>2</sub> CF <sub>3</sub>	460	1A
	-SO <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	434	1B
	-SO <sub>2</sub> CH <sub>3</sub>	456	1C
	-SO <sub>2</sub> CH <sub>3</sub>	456	1D
	-SO <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	484	1E
	-SO <sub>2</sub> CF <sub>3</sub>	510	1F
	-SO <sub>2</sub> CH <sub>3</sub>	472	1G
	-SO <sub>2</sub> CH <sub>3</sub>	406	1H




Y	R <sup>b</sup>	MS (M+H)	Example
	-SO <sub>2</sub> CF <sub>3</sub>	460	1I
	-SO <sub>2</sub> CH <sub>3</sub>	456	1J
	-SO <sub>2</sub> CH <sub>3</sub>	424	1K

Example 25 Step 1

A stirred solution of 1M 1-thienyllithium in THF (40 ml, 40 mmol) was cooled in a dry-ice/acetone bath under N<sub>2</sub>. Triethylborate (8.5 ml, 50 mmol) was added, and the reaction mixture was allowed to warm to R.T.. After 20 min., 4-iodoaniline (6.6 g, 30 mmol), Na<sub>2</sub>CO<sub>3</sub> (4.5 g), H<sub>2</sub>O (20 ml), and Pd(dppf)Cl<sub>2</sub>·CH<sub>2</sub>Cl<sub>2</sub> (750 mg, 0.9 mmol) were added. The reaction mixture was stirred under N<sub>2</sub> until the exotherm was complete, then partitioned between Et<sub>2</sub>O and H<sub>2</sub>O. The Et<sub>2</sub>O layer was washed with 1N NaOH, dried (Na<sub>2</sub>CO<sub>3</sub>), and filtered through a pad of silica gel, eluting with Et<sub>2</sub>O. The resultant brown solid was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 ml) and a solution of trifluoroacetic anhydride (8 ml, 57 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 ml) was added in portions with stirring. To the resultant suspension was added CH<sub>2</sub>Cl<sub>2</sub> (450 ml) and the reaction mixture was stirred for 20 min. Water (200 ml) was added, followed by NaHCO<sub>3</sub> (7 g) in portions until CO<sub>2</sub> evolution ceased. The organic layer was stirred with MgSO<sub>4</sub> and DARCO, then filtered and concentrated to give a solid. The solid was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 ml) and to the stirred solution was added hexanes (100 ml). The solid was collected, washed with hexanes and dried to give the product (6.12 g, 75%). M.p.


213-216 °C. Calcd for  $C_{12}H_8F_3NOS$ : C, 53.14; H, 2.58; N, 5.17. Found: C, 53.06; H, 2.85; N, 4.90%.

### Step 2




To a solution of the product of Step 1 (19.0 g, 70 mmol) in DMF (150 ml) was added N-chlorosuccinimide (10.1 g, 76 mmol) and trifluoroacetic acid (1.5 ml), and the reaction mixture was stirred under  $N_2$  for 2 days. Water (500 ml) was added and the resultant solid was collected, washed with water and dried to give the product (20.6 g, 96%). M.P. 198 - 200 °C. Calcd for  $C_{12}H_7ClF_3NOS$ : C, 47.12; H, 2.29; N, 4.58. Found: C, 47.19; H, 2.15; N, 4.47%.

### Step 3



A mixture of the product of Step 2 (15.0 g, 49.1 mmol) and sodium hydroxide (19.6 g, 490 mmol) in MeOH (400 ml) and water (150 ml) was stirred at R.T. overnight. The mixture was concentrated *in vacuo* and the residue was partitioned between EtOAc and water. The organic layer was washed with water, brine, dried, and concentrated. The residue was purified by flash column (1:3 acetone/hexanes) to give the product (10.14 g, 98%).  $^1H$ -NMR ( $CDCl_3$ , 400 MHz)  $\delta$  7.32 (2H, m), 6.90 (1H, d,  $J=4.8$  Hz), 6.83 (1H, d,  $J=4.8$  Hz), 6.67 (2H, m), 3.76 (2H, b).

### Step 4




To a stirred, ice-cold solution of the product of Step 3 (2.0 g, 9.5 mmol) in THF (100 ml) was added pyridine (2.3 ml, 28 mmol) and N,N'-disuccinimidyl carbonate

26

(2.44 g, 9.5 mmol). The reaction mixture was stirred at ice-bath temp. for 1.5 h, then Preparation 1 (2.04 g, 9.5 mmol) was added, and the reaction mixture was allowed to warm to R.T.. After 16 h, the reaction mixture was concentrated, the residue was dissolved in EtOAc (200 ml) and washed with 2N HCl, sat'd NaHCO<sub>3</sub> and sat'd NaCl.

- 5 The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and evaporated to afford the product (4.21 g, 98%) that was used directly in Step 5. HRMS calc. for C<sub>22</sub>H<sub>29</sub>ClN<sub>3</sub>O<sub>3</sub>S (M+H) 450.1618. Found 450.1623.

#### Step 5




Reaction of the product of Step 4 (4.11 g, 9.13 mmol) with HCl by the procedure of Example 1, Step 3 afforded the product (3.71 g) that was used directly in Step 6. HRMS calc. for C<sub>17</sub>H<sub>21</sub>ClN<sub>3</sub>O<sub>3</sub>S (M+H) 350.1094. Found 350.1100.

15 


#### Step 6

- To a suspension of the product of Step 5 (50 mg, 0.13 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was added Et<sub>3</sub>N (39 mg, 0.39 mmol) followed by n-propylsulfonyl chloride (20 mg, 0.14 mmol). The reaction mixture was stirred for 16 h. EtOAc (10 ml) was added and the mixture was washed with 2N HCl, sat'd NaHCO<sub>3</sub> and sat'd NaCl, dried (MgSO<sub>4</sub>),  
20 filtered and concentrated. The residue was subjected to PTLC (3:97 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (37 mg, 62%). HRMS calc. for C<sub>20</sub>H<sub>27</sub>ClN<sub>3</sub>O<sub>3</sub>S<sub>2</sub> (M+H) 456.1182. Found 456.1179.


Reaction of the product of Step 5, 2-5-1, with the appropriate sulfonyl chloride in the presence of Et<sub>3</sub>N gave the following examples.



R <sup>6</sup>	MS (M+H) <sup>+</sup>	Example
-SO <sub>2</sub> CH <sub>3</sub>	428	2A
-SO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	442	2B
-SO <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	456	2C
-SO <sub>2</sub> CF <sub>3</sub>	482	2D
-SO <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	496	2E


Example 3

5

Step 1

Using the procedure of Example 1, Step 1, Preparation 1 (2.3 g, 107 mmol) was reacted with 4-iodophenyl isocyanate (2.6 g, 107 mmol). Purification by flash chromatography (2:98 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) afforded a white solid.


10

Step 2

A mixture of the product of Step 1 (3.0 g, 6.7 mmol), 4M HCl in 1,4-dioxane (15 ml) and THF (15 ml) was stirred at ambient temp. for 5 h. The reaction mixture was concentrated to dryness, and H<sub>2</sub>O (100 ml) and 3M NaOH (20 ml) was added to the residue. The whole was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x100 ml). The combined organic extracts were dried (MgSO<sub>4</sub>), filtered and evaporated. Flash chromatography (2:98 MeOH/CH<sub>2</sub>Cl<sub>2</sub> then 10:90 (2M NH<sub>3</sub> in MeOH)/CH<sub>2</sub>Cl<sub>2</sub>) gave a white solid (2.4 g, 100%). HRMS calc. for C<sub>13</sub>H<sub>19</sub>IN<sub>3</sub>O (M+H) 360.0573. Found 360.0576.

15

20


Step 3

- To a stirred ice-cold mixture of the product of Step 2 (2.4 g, 6.7 mmol) and cyclopropane carboxaldehyde (0.8 ml, 11 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 ml) was added
- 5  $\text{NaBH}(\text{OAc})_3$  (1.83 g, 10.8 mmol). The reaction mixture was allowed to warm to room temp. and stirred overnight. The reaction mixture was cooled in ice and 3M NaOH (5 ml) was added. After 0.5 h the mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3x100 ml), dried ( $\text{MgSO}_4$ ), filtered and evaporated. The residue was triturated with  $\text{CH}_2\text{Cl}_2$ /hexanes (1:10) to afford a white solid (2.4 g, 87%). HRMS calc. for  $\text{C}_{17}\text{H}_{25}\text{IN}_3\text{O}$  (M+H)
- 10 414.1038. Found: 414.1042.

Step 4




- A vessel charged with the product of Step 3 (200 mg, 0.48 mmol), 4-trifluoromethoxybenzeneboronic acid (250 mg, 1.21 mmol),
- 15 tris(dibenzylideneacetone)dipalladium (0) (50 mg, 0.05 mmol),  $\text{CsCO}_3$  (0.8 g, 2.5 mmol) and toluene (10 ml) was refluxed under  $\text{N}_2$  for 3 h. The reaction mixture was allowed to cool, then EtOAc (50 ml) and  $\text{H}_2\text{O}$  (25 ml) were added. Solids were removed by filtration and the EtOAc layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and evaporated. The residue was subjected to PTLC (3:7 acetone/hexanes then 10:90
- 20 (2M  $\text{NH}_3$  in MeOH)/ $\text{CH}_2\text{Cl}_2$ ) to give a pale yellow solid (50 mg, 23%). HRMS calc. for  $\text{C}_{24}\text{H}_{29}\text{F}_3\text{N}_3\text{O}_2$  (M+H) 448.2212. Found 448.2215.



Using appropriate starting materials and essentially the same procedure, the following compounds were prepared:



Y	MS (M+H) <sup>+</sup>	Example
	364.1	3A
	382	3B
	404	3C

29


Y	MS (M+H) <sup>+</sup>	Example
	423	3D
	400	3E
	382	3F

Example 4Step 1


To an N<sub>2</sub>-purged mixture of 4-bromonitrobenzene (20.0 g, 99.0 mmol), 3,5-difluorophenylboronic acid (23.4 g, 148 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (38.7 g, 119 mmol) in toluene (600 ml) and H<sub>2</sub>O (30 ml) was added Pd(dppf)Cl<sub>2</sub>·CH<sub>2</sub>Cl<sub>2</sub> (4.04 g, 4.95 mmol). The reaction mixture was heated at 90 °C for 2 h, allowed to cool to R.T., then filtered through celite. The whole was extracted with EtOAc (3x500 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to give a solid. To a vigorously stirred ice-cold mixture of the solid in CH<sub>3</sub>OH (1 L) and NiCl<sub>2</sub>·6H<sub>2</sub>O (61.0 g, 257 mmol) was added NaBH<sub>4</sub> (14 g, 370 mmol) in portions. After the addition was complete, the reaction mixture was poured into H<sub>2</sub>O (100 ml), then filtered through celite and extracted with EtOAc (3x500 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The residue was dissolved in EtOAc, and 1N HCl/Et<sub>2</sub>O (300 ml) was added. The precipitate was washed with hexane, air-dried, and dissolved in H<sub>2</sub>O. The solution was neutralized by addition of 1N NaOH, then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x1 L). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to give the product (19.0 g, 94%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400

MHz)  $\delta$  7.38 (2H, m), 7.06 (2H, m), 6.75 (2H, m), 6.72 (1H, m), 3.81 (s, 2H). MS m/e 206 (M+H).

Using the appropriate substituted phenylboronic acid starting material and essentially the same procedure, the following compounds were prepared:




$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.41-7.21 (5H, m), 7.33 (1H, m), 6.76 (2H, m), 3.76 (2H, b).



$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.39 (2H, m), 7.24 (3H, m), 6.76 (2H, m), 3.80 (2H, b).


10 Additional arylamines were prepared from 4-iodoaniline according to the following procedure.




15 A mixture of 4-iodoaniline (1.00 g, 4.57 mmol), 3-trifluoromethylphenylboronic acid (1.30 g, 6.85 mmol) and  $\text{Cs}_2\text{CO}_3$  (1.64 g, 5.02 mmol) in toluene (50 ml) and  $\text{H}_2\text{O}$  (3 ml) was purged with  $\text{N}_2$  for 5 min. To the reaction mixture was added  $\text{Pd}(\text{dppf})\text{Cl}_2 \cdot \text{CH}_2\text{Cl}_2$  (746 mg, 0.91 mmol). The reaction mixture was heated at 90 °C for 5 h, then allowed to cool to R.T. and poured into cold water. The whole was extracted with  $\text{CH}_2\text{Cl}_2$  (3x100 ml). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. Purification of the residue by PTLC (EtOAc/hexane 1:2) gave

20 the product (216 mg, 20%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.77 (1H, m), 7.70 (1H, m), 7.51 (2H, m), 7.42 (2H, m), 6.78 (2H, m), 3.65 (2H, b).

Using the appropriate substituted phenylboronic acid starting material and essentially the same procedure, the following compounds were prepared.




$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.54 (1H, m), 7.34 (3H, m), 7.15 (1H, t,  $J = 8.8$  Hz), 6.75 (2H, m), 3.76 (2H, b).



4-1-6

- 5  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.48 (2H, m), 7.35 (2H, d,  $J = 6.4$  Hz), 7.08 (2H, t,  $J = 6.4$  Hz), 6.76 (2H, d,  $J = 6.4$  Hz), 3.73 (2H, b). MS  $m/e$  188 ( $\text{M}+\text{H}$ ).




4-1-7

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.51 (1H, m), 7.41 (3H, m), 7.32 (1H, m), 7.23 (1H, m), 6.75 (2H, m), 3.78 (2H, b). MS  $m/e$  204 ( $\text{M}+\text{H}$ ).

10


### Step 2



4-2-1

- A stream of  $\text{N}_2$  was passed through a mixture of the product of Preparation 2 (2.00 g, 9.33 mmol), 3-bromopyridine (2.95 g, 18.7 mmol) and 2-(di-*tert*-butylphosphino)biphenyl (0.139 g, 0.467 mmol) and NaOtBu (1.80 g, 18.7 mmol) in anhydrous toluene (10 ml).  $\text{Pd}(\text{OAc})_2$  (0.105 g, 0.467 mmol) was added and the reaction mixture was stirred at 110 °C for 24 h. The reaction mixture was allowed to cool to R.T. and poured into cold  $\text{H}_2\text{O}$ . The whole was extracted with  $\text{CH}_2\text{Cl}_2$  (3x50 ml) and the combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated. Purification of the residue by PTLC (1:20  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) gave the product (1.47 g, 54%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  8.29 (1H, s), 8.07 (1H, b), 7.17 (2H, m), 4.2 (1H, b), 3.74 (2H, m), 2.82 (2H, m), 2.74 (3H, s), 1.70 (4H, m), 1.45 (9H, s). MS  $m/e$  292 ( $\text{M}+\text{H}$ ).




Step 3

4-3-1


To the product of Step 2 (1.47 g, 5.05 mmol) was added 4M HCl/1,4-dioxane (20 ml). The reaction mixture was stirred at R.T. for 1.5 h and concentrated to afford the product in quantitative yield. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 400 MHz) δ 8.46 (1H, s), 8.14 (2H, m), 7.86 (1H, s), 4.13 (2H, m), 3.40 (1H, b), 3.16 (2H, b), 2.75 (3H, s), 2.26 (2H, m), 1.76 (2H, m). MS m/e 192 (M+H).

Step 4

To a mixture of the product of Step 1 (4-1-1) (0.100 g, 0.487 mmol) and *i*Pr<sub>2</sub>NEt (0.43 ml, 2.44 mmol) in anhydrous toluene (10 ml) was added triphosgene (0.051 g, 0.171 mmol). The mixture was stirred at 120 °C for 2 h, then allowed to cool to R.T., and the product of Step 3 (4-3-1) (0.133 g, 0.585 mmol) was added. The reaction mixture was stirred at R.T. for 16 h, then poured into cold H<sub>2</sub>O and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x20 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The residue was purified by PTLC (1:20 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (0.114 g, 56 %). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.33 (1H, d, *J* = 2.4 Hz), 8.09 (1H, m), 7.49 (4H, m), 7.17 (2H, m), 7.06 (2H, m), 6.74 (1H, m), 6.51 (1H, s), 4.49 (1H, m), 3.77 (2H, m), 2.93 (3H, s), 2.91 (2H, m), 1.85 (4H, m). MS m/e 423 (M+H).

Example 5

5


Step 1

5-1-1

The product 5-1-1 was prepared in 57% yield from 2-bromopyridine and Preparation 2 by the procedure of Example 4, Step 2, except that 2-(di-tert-

butylphosphino)biphenyl was replaced by 1,3-bis(diphenylphosphino)propane, and a reaction temperature of 80 °C instead of 110 °C was used. MS m/e 292 (M+H).

### Step 2




5-2-1

Treatment of the product of Step 1 with 4 N HCl/dioxane by the procedure of Example 4, Step 3 gave the product. MS m/e 192 (M+H).

### Step 3

To a stirred ice-cold mixture of 4-1-2 (0.063 g, 0.339 mmol) and pyridine (0.14 ml, 1.69 mmol) in anhydrous THF (10 ml) was added N,N'-disuccinimidyl carbonate (0.087 g, 0.339 mmol). The reaction was stirred in an ice-bath for 25 min. then the product of Step 2, 5-2-1 (0.100 g, 0.508 mmol), was added. The reaction was allowed to warm to R.T., stirred for 16 h, then poured into cold H<sub>2</sub>O (20 ml). The whole was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x20 ml), the combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The residue was subjected to PTLC (1:20 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (0.080 g, 58%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.19 (1H, m), 7.52 (5H, m), 7.37 (2H, m), 7.27 (1H, m), 6.99 (1H, m), 6.69 (1H, d), 6.62 (1H, m), 6.45 (1H, s), 4.56 (1H, m), 4.42 (2H, m), 2.92 (2H, m), 2.88 (3H, s), 1.78 (4H, m). MS m/e 405 (M+H).


### Example 6



Reaction of 4-1-4, N,N'-disuccinimidyl carbonate and 5-2-1 by the procedure of Example 5, Step 3 afforded the product. MS m/e 455 (M+H).


MISSING AT THE TIME OF PUBLICATION

35


Example 11

Reaction of 4-1-2, triphosgene and 4-3-1 by the procedure of Example 4, Step 4 afforded the product. MS m/e 405 (M+H).


5

Example 12

10 Reaction of 4-1-7, triphosgene and 4-3-1 by the procedure of Example 4, Step 4 afforded the product. MS m/e 421 (M+H).


Example 13

15

Step 1

A mixture of Preparation 3 (2.75 g, 9.7 mmol), 2-bromothiazole (1.98 g, 12.1 mmol), and K<sub>2</sub>CO<sub>3</sub> (3.5 g, 25 mmol) in DMF (40 ml) was heated at 160 °C for 20 h.

20 The reaction mixture was concentrated and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The organic layer was washed with sat'd NaCl, dried (MgSO<sub>4</sub>), filtered and concentrated. Flash chromatography (gradient; CH<sub>2</sub>Cl<sub>2</sub> to 2:98 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave the product (2.0 g, 62%). MS m/e 332.1 (M+H).


Step 2

13-2-1

The product of Step 1 (2.0 g, 6.0 mmol) and 33% HBr in AcOH (40 ml) was stirred at R.T. for 2 h. The reaction mixture was evaporated and the residue was partitioned between 1N NaOH and CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with sat'd NaCl, dried (MgSO<sub>4</sub>), filtered and evaporated. Flash chromatography (gradient; 2:98 (2M NH<sub>3</sub> in MeOH)/CH<sub>2</sub>Cl<sub>2</sub> to 15:85 (2M NH<sub>3</sub> in MeOH)/CH<sub>2</sub>Cl<sub>2</sub>) gave the product (0.94 g, 79%) as a yellow solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.04 (1H, d, J = 4 Hz), 6.52 (1H, d, J = 4 Hz), 3.96 (2H, m), 3.17 (1H, m), 2.99 (2H, m), 2.59 (3H, s), 2.16 (2H, m), 1.68 (2H, m). MS m/e 198 (M+H).


Step 3

Reaction of 4-1-2, triphosgene and 13-2-1 by the procedure of Example 4, Step 4 afforded the product. MS m/e 411 (M+H).


Example 14

14

Reaction of 4-1-1, triphosgene and 13-2-1 by the procedure of Example 4, Step 4 afforded the product. MS m/e 429 (M+H).


Example 15

15

Step 1

15-1-1

An N<sub>2</sub>-purged mixture of 2-bromopyrimidine (400 mg, 2.52 mmol), Preparation 3 (510 mg, 1.79 mmol), Pd(OAc)<sub>2</sub> (18 mg, 0.08 mmol), sodium *tert*-butoxide (516 mg, 5.37 mmol), and (1,3-bis—diphenylphosphino)propane (29 mg, 0.07 mmol) in toluene (6 ml) was stirred at 70 °C in a sealed vessel for 16 h. The reaction mixture was allowed to cool to R.T., and 1N NaOH (20 ml) was added. The whole was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x20 ml), and the combined CH<sub>2</sub>Cl<sub>2</sub> extracts were dried (MgSO<sub>4</sub>), filtered, and evaporated. The residue was subjected to PTLC (2:98 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (464 mg, 79%). MS m/e 327 (M+H).

Step 2


15-2-1

The product of Step 1 (464 mg, 1.43 mmol) and 10% Pd/C (59 mg) in EtOH (20 ml) was stirred under 1 atm. of H<sub>2</sub> for 16 h. The catalyst was removed by filtration through celite and the filter pad was washed with EtOH. The combined filtrate and washings were evaporated. The residue was subjected to PTLC (5:95 (2M NH<sub>3</sub> in MeOH)/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (464 mg, 79%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.28 (2H, m), 6.44 (1H, m), 4.66 (2H, m), 2.99 (2H, m), 2.65 (1H, m), 2.47 (3H, s), 1.96 (2H, m), 1.33 (2H, m). MS m/e 193 (M+H).



Step 3

Reaction of the product of Step 2 (15-2-1) with 4-1-2 with triphosgene by the procedure of Example 4, Step 4 gave the product. MS (m/e) 406 (M+H).

38

Example 16


Reaction of the product of Example 15, Step 2 (15-2-1) and 4-1-1 with  
 5 triphosgene by the procedure of Example 4, Step 4 gave the product. MS (m/e) 424 (M+H).


Example 1710 Step 1

Reaction of the product of Example 5, Step 2 with  
 4-bromo-2-fluorophenylisocyanate by the procedure of Example 1, Step 1 gave the  
 product. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 8.18 (1H, m), 7.47 (1H, m), 7.38 (2H, m), 7.30  
 15 (2H, m), 6.68 (1H, m), 6.61 (1H, m), 4.49 (1H, m), 4.43 (2H, m), 2.91 (2H, m), 2.85  
 (3H, s), 1.71 (4H, m). MS m/e 391 (M+H).

Step 2


Reaction of the product of Step 1 with 3-fluorophenylboronic acid by the  
 20 procedure of Example 4, Step 1 gave the product. MS m/e 423 (M+H).

Example 18

Step 1

5 A mixture of 4-biphenyl isocyanate (3.00 g, 15.4 mmol) and Preparation 1 (5.33 g, 25.0 mmol) in  $\text{CH}_2\text{Cl}_2$  (100 ml) was stirred at R.T. for 16 h. The mixture was washed with water (25 ml), 3N HCl (25 ml), and brine (50 ml). The organic portion was dried ( $\text{MgSO}_4$ ), filtered, concentrated, and purified by column chromatography (gradient;  $\text{CH}_2\text{Cl}_2$  to 1:99  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) to give the product (6.11 g, 97%). MS (ES) m/e 410 ( $\text{M}+\text{H}$ )<sup>+</sup>.

10

Step 2

15 A mixture of the product of Step 1 (6.11 g, 14.9 mmol) and 4N HCl/dioxane (100 ml) was stirred at R.T. for 5 h. The volatiles were evaporated and the residue was triturated with ether. The precipitate was collected, dissolved in water (200 ml), basified to pH 14, and extracted with  $\text{CH}_2\text{Cl}_2$  (300 ml). The organic portion was dried and concentrated to give the product (4.39 g, 92%). MS (ES) m/e 310 ( $\text{M}+\text{H}$ )<sup>+</sup>.

20 Step 3

A solution of the product of Step 2 (80 mg, 0.26 mmol), nicotinoyl chloride hydrochloride (54 mg, 0.30 mmol), and triethylamine (90  $\mu\text{l}$ , 0.64 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 ml) was stirred at R.T. for 16 h. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (50 ml) and extracted with 3N NaOH (5 ml). The organic layer was washed with water (15 ml), dried, ( $\text{MgSO}_4$ ), filtered, and concentrated. The residue was subjected to PTLC (4:96  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) to give the product (90 mg, 84%). <sup>1</sup>H NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  8.68 (2H, m), 7.76 (1H, m), 7.2-7.6 (10H, m), 6.48 (1H, s), 4.85 (1H, m),




40

4.60 (1H, m), 3.80 (1H, m), 3.20 (1H, m), 2.91 (3H, s), 2.86 (1H, m), 1.4-2.0 (4H, m).

MS (ES)  $m/e$  415 ( $M+H$ )<sup>+</sup>.

Using the appropriate acid chloride and essentially the same procedure the following compounds were prepared.




5

R <sup>6</sup>	(M+H) <sup>+</sup>	Example
C(O)CH <sub>3</sub>	352	18B
C(O)-	378	18C
C(O)-	420	18D
C(O)-	414	18E
C(O)-	415	18F
C(O)-	415	18G

#### Example 19





Reaction of Example 1, 1-3-5, with the appropriate acid chloride afforded the following compounds:



10


R <sup>6</sup>	(M+H) <sup>+</sup>	Example
C(O)-CH <sub>3</sub>	370	19A
C(O)-	396	19B
C(O)-	432	19C
C(O)-	433	19D
C(O)-	433	19E
C(O)-	433	19F

41






R <sup>6</sup>	(M+H) <sup>+</sup>	Example
	467	19G
	501	19H
	481	19I
	497	19J

Example 20

Reaction of the product of Example 1, 1-3-7, with the appropriate acid chloride afforded the following compounds:




5






R	(M+H) <sup>+</sup>	Example
C(O)-CH <sub>3</sub>	388	20A
C(O)- 	414	20B
C(O)- 	450	20C
C(O)- 	451	20D
C(O)- 	451	20E
C(O)- 	451	20F


Example 21

Reaction of the product of Example 2, Step 5, 2-5-1, with the appropriate acid chloride afforded the following compounds




10


R <sup>6</sup>	(M+H) <sup>+</sup>	Example
C(O)-CH <sub>3</sub>	392	21A
C(O)- 	418	21B
C(O)- 	454	21C
C(O)- 	455	21D
C(O)- 	455	21E
C(O)- 	455	21F

Example 22

A mixture of Example 18 (45 mg, 0.11 mmol) and 3-chloroperoxybenzoic acid  
 5 (40 mg) in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) was stirred at R.T. for 16 h. The mixture was diluted with  
 CH<sub>2</sub>Cl<sub>2</sub> (50 ml), then washed with 3N NaOH (2x5 ml) and water (10 ml). The organic  
 layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated. The residue was subjected to  
 PTLC (1:9 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (34 mg, 73%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400  
 MHz) δ 8.20 (2H, m), 7.2-7.6 (11H, m), 6.56 (1H, s), 4.76 (1H, m), 4.59 (1H, m), 3.78  
 10 (1H, m), 3.22 (1H, m), 2.7-3.0 (4H, m), 1.4-2.0 (4H, m). MS (ES) m/e 431 (M+H)<sup>+</sup>.

Example 23


23

Step 1

15


A mixture of 4-piperidone ethylene ketal (0.64 ml, 5.0 mmol) and sulfamide (0.53 g, 5.5 mmol) in DME (20 ml) was refluxed for 16 h. The mixture was concentrated to ca. 3 ml, dissolved in EtOAc (175 ml), washed with sat'd  $\text{NH}_4\text{Cl}$  (2x25 ml), water (2x25 ml), and brine (25 ml). The organic portion was dried, filtered, and evaporated to give the product (0.58 g, 52%). MS (ES)  $m/e$  223 ( $\text{M}+\text{H}$ )<sup>+</sup>.

### Step 2



A mixture of the product of Step 1 (560 mg, 2.52 mmol) and pyridinium 4-toluenesulfonate (190 mg, 0.756 mmol) in acetone (25 ml) and water (0.5 ml) was refluxed for 64 h. The mixture was evaporated to dryness and the residue was partitioned between  $\text{CH}_2\text{Cl}_2$  (75 ml) and aq.  $\text{NaHCO}_3$  (2x20 ml). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  and EtOAc sequentially. The EtOAc layer was evaporated to give the product (140 mg). <sup>1</sup>H NMR ( $\text{CD}_3\text{OD}$ , 400 MHz)  $\delta$  3.47 (1H, t,  $J=6.4$  Hz), 3.15 (3H, m), 2.54 (1H, t,  $J=6.4$  Hz), 1.81 (3H, m).

### Step 3



A mixture of the product of Step 2 (135 mg, 0.757 mmol), 40% aqueous methylamine (300  $\mu\text{l}$ , 2.42 mmol), and sodium triacetoxyborohydride (375 mg, 1.77 mmol) in dichloroethane (5 ml) was stirred at R.T. for 19 h. The mixture was partitioned between 3N NaOH (5 ml) and EtOAc (3x50 ml). The organic layer was concentrated to give the crude product (40 mg). The aqueous layer was evaporated *in vacuo* to dryness and the residue was suspended in EtOAc. The suspension was filtered and the filtrate concentrated to give another batch of the product (70 mg). MS (FAB)  $m/e$  194 ( $\text{M}+\text{H}$ )<sup>+</sup>.


Step 4

To an ice-cold solution of 4-1-2 (40 mg, 0.21 mmol) in anhydrous THF (3 ml) was added N,N'-disuccinimidyl carbonate (55 mg, 0.21 mmol) and pyridine (52  $\mu$ l, 0.65 mmol). The mixture was stirred at 0 °C for 2 h and the product of Step 3 (70 mg, 0.36 mmol) was added. After stirring at R.T. for 2 h the reaction mixture was taken up in CH<sub>2</sub>Cl<sub>2</sub> (50 ml), washed with 1N HCl (10 ml), dried, (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The residue was subjected to PTLC (5:95 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (62 mg, 71%).


- <sup>1</sup>H NMR (CD<sub>3</sub>OD, 400 MHz)  $\delta$  7.56 (2H, m), 7.48 (2H, m), 7.40 (2H, m), 7.32 (1H, m), 7.02 (1H, m), 4.23 (1H, m), 3.75 (2H, m), 2.94 (3H, s), 2.72 (2H, m), 1.7-2.0 (4H, m). MS (ES) m/e 407 (M+H)<sup>+</sup>.

Using the appropriate starting materials and essentially the same procedure afforded the following compounds.

15



Y	(M+H) <sup>+</sup>	Example
	389	23A
	425	23B
	429	23C

Example 24


24

- A mixture of 1-3-5 (71 mg, 0.20 mmol), 2-bromoacetamide (32 mg, 0.23 mmol), and anhydrous potassium carbonate (170 mg, 1.20 mmol) in CH<sub>3</sub>CN (2 ml) in a


45

sealed tube was heated to 45 °C for 6 h. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (75 ml), washed with water (50 ml), dried, and concentrated. The residue was subjected to PTLC (5:95 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (37 mg, 49%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.48 (4H, m), 7.35 (2H, m), 7.23 (1H, m), 6.98 (2H, m), 6.56 (1H, s), 5.97 (1H, bs), 4.25 (1H, m), 2.8-3.0 (7H, m), 2.31 (2H, m), 1.6-1.8 (4H, m). MS (ES) m/e 385 (M+H)<sup>+</sup>.

### Example 25




25A



25B

10


#### Step 1



25-1-1

To ethyl 4-oxocyclohexanecarboxylate (10 g, 59 mmol) in MeOH (75 ml) and water (50 ml) was added lithium hydroxide monohydrate (4.2 g, 100 mmol) at 0 °C. The mixture was warmed up to R.T. and stirred for 3 h. The mixture was acidified to pH 2 with 3N HCl. The volatiles were evaporated and the residue was extracted with EtOAc (300 ml). The organic portion was dried and concentrated to give the product (8.01 g, 96%). MS (CI) m/e 143 (M+H)<sup>+</sup>.

#### Step 2




25-2-1

2M oxalyl chloride in CH<sub>2</sub>Cl<sub>2</sub> (20 ml, 40 mmol) was added over 5 min to a solution of the product of Step 1 (3.0 g, 21 mmol) in anhydrous THF (50 ml). The solution was heated to 80 °C for 6 h and then evaporated to dryness. The residue was dissolved in THF (50 ml) at 0 °C and aq. NH<sub>4</sub>OH (6.0 ml, 89 mmol) was added. After stirring at R.T. for 16 h, the mixture was concentrated and the residue purified by

column chromatography (gradient  $\text{CH}_2\text{Cl}_2$  to 2:98  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) to give the product (762 mg, 26%). MS (CI) m/e 142 ( $\text{M}+\text{H}$ )<sup>+</sup>.

### Step 3



A mixture of the product of Step 2 (800 mg, 5.71 mmol), 40% aq. methylamine (4.0 ml, 52 mmol), and sodium triacetoxyborohydride (1.7 g, 8.0 mmol) in dichloroethane (20 ml) was stirred at R.T. for 16 h. The reaction was quenched with 3N NaOH and partitioned between brine and 1:1  $\text{CH}_3\text{CN}/\text{CH}_2\text{Cl}_2$ . The organic portion  
10 was concentrated and the residue purified by column chromatography (gradient  $\text{CH}_2\text{Cl}_2$  to 1:4 2M  $\text{NH}_3$  in  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) to give the product (450 mg, 51%). MS (CI) m/e 157 ( $\text{M}+\text{H}$ )<sup>+</sup>.

### Step 4

15 A mixture of the aniline 4-1-2 (100 mg, 0.534 mmol), N,N'-disuccinimidyl carbonate (137 mg, 0.535 mmol), and pyridine (0.13 ml, 1.6 mmol) in THF (3 ml) was stirred at 0 °C for 2 h. To this mixture was added the product of Step 3 (125 mg, 0.811 mmol) and the reaction was stirred at R.T. for 2 h. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (100 ml), washed with 1N HCl (2x25 ml), water (2x25 ml), brine (25 ml), dried,  
20 and concentrated. The residue was subjected to PTLC (3:97  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) to give the *cis*-product (14 mg) and the *trans*-product (15 mg).

#### *cis*-product 25A:

<sup>1</sup>H NMR ( $\text{CD}_3\text{OD}$ , 400 MHz):  $\delta$  7.4-7.6 (4H, m), 7.33 (2H, m), 7.22 (1H, m), 6.95 (1H, m), 4.13 (1H, m), 2.86 (3H, s), 2.53 (1H, m), 2.13 (2H, m), 1.82 (2H, m), 1.5-1.75 (4H, m). MS (ES) m/e 370 ( $\text{M}+\text{H}$ )<sup>+</sup>.


25

#### *trans*-product 25B:


47

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 400 MHz):  $\delta$  7.4-7.5 (4H, m), 7.34 (2H, m), 7.23 (1H, m), 6.96 (1H, m), 4.07 (1H, m), 2.88 (3H, s), 2.14 (1H, m), 1.98 (2H, m), 1.81 (2H, m), 1.5-1.7 (4H, m). MS (ES)  $m/e$  370 ( $\text{M}+\text{H}$ ) $^+$ .

- 5 Reaction of the product of Step 3, 25-3-1 with aniline 4-1-1 by essentially the same procedure gave 25C and 25D:




25C MS (ES)  $m/e$  388 ( $\text{M}+\text{H}$ ) $^+$



25D MS(ES)  $m/e$  388 ( $\text{M}+\text{H}$ ) $^+$


10

### Example 26



26

### Step 1



15

26-1-1


- To a stirred mixture of 1,4-cyclohexanedione monoethylene ketal (4.68 g, 30 mmol) and 40% w/w aq. methylamine (6.0 mL) in 1,2-dichloroethane (75 mL), was added  $\text{Na}(\text{OAc})_3\text{BH}$  (9.6 g, 45 mmol) in portions. The reaction mixture was vigorously stirred for 16 h, then 1N NaOH (75 mL) was added. The organic layer was washed with sat'd NaCl, dried ( $\text{MgSO}_4$ ), filtered, and evaporated to give an oil (4.60 g, 90%)
- 20



48

that was used without further purification.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  3.97 (4H, s), 2.47 (1H, m), 2.46 (3H, s), 1.91 (2H, m), 1.80 (2H, m), 1.59 (2H, m), 1.45 (2H, m).

### Step 2




To a stirred, ice-cold mixture of aniline 4-1-1 (1.00 g, 4.87 mmol) and pyridine (1.97 ml, 24.3 mmol) in anhydrous THF (50 ml) was added disuccinimidyl carbonate (1.25 g, 4.87 mmol). The reaction mixture was stirred at 0 °C for 1 h and the product of Step 1 (1.25 g, 7.31 mmol) was added. The reaction mixture was allowed to warm to R.T., stirred for 16 h, then poured into cold  $\text{H}_2\text{O}$  (100 ml). The whole was extracted with  $\text{CH}_2\text{Cl}_2$  (3x100 ml). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and evaporated. Purification of the residue by column chromatography (1:20  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) afforded the product (1.40 g, 71%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.49 (4H, m), 7.10 (2H, m), 6.70 (1H, m), 6.60 (1H, s), 4.30 (1H, m), 3.90 (4H, s), 2.90 (3H, s), 1.75 (8H, m). MS m/e 403 (M+H).

10

15

### Step 3





To the product of Step 2 (1.30 g, 3.23 mmol) in THF (30 ml) was added 5N HCl (20 ml). The reaction mixture was stirred at R.T. for 4.5 h, then extracted with  $\text{CH}_2\text{Cl}_2$  (3x100 ml). The combined organic extracts were washed with sat'd  $\text{NaHCO}_3$ , dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. The residue was purified by PTLC (1:20  $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ ) to give the product (0.80 g, 69%).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400 MHz)  $\delta$  7.50

20


(4H, m), 7.10 (2H, m), 6.80 (1H, m), 6.50 (1H, s), 4.80 (1H, m), 2.90 (3H, s), 2.48 (4H, m), 2.10 (2H, m), 1.90 (2H, m). MS m/e 359 (M+H).

#### Step 4



To a mixture of the product of Step 3 (0.43 g, 1.20 mmol) and benzylamine (0.257 g, 2.40 mmol) in 1,2-dichloroethane (10 ml) was added NaBH(OAc)<sub>3</sub> (0.762 g, 3.60 mmol) in portions. The reaction mixture was stirred at R.T. for 4.5 h, then poured into sat'd NaHCO<sub>3</sub> (20 ml) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x20 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated. The residue was purified by PTLC (1:20 (2M NH<sub>3</sub>/CH<sub>3</sub>OH):CH<sub>2</sub>Cl<sub>2</sub>) to produce the *cis*-isomer 26-4-1 (0.240 g, 44.5%) and the *trans*-isomer 26-4-2 (0.200 g, 37.0%). *Cis* isomer: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.48 (4H, m), 7.30 (5H, m), 7.05 (2H, m), 6.70 (1H, m), 6.40 (1H, s), 4.20 (1H, m), 3.78 (2H, s), 2.90 (4H, m), 1.90 (4H, m), 1.55 (4H, m). MS m/e 450 (M+H). *Trans*-isomer: <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.48 (4H, m), 7.33 (5H, m), 7.05 (2H, m), 6.70 (1H, m), 6.37 (1H, s), 4.20 (1H, m), 3.82 (2H, s), 2.88 (3H, m), 2.50 (1H, m), 2.10 (2H, m), 1.80 (2H, m), 1.20-1.70 (4H, m). MS m/e 450 (M+H).


#### 20 Step 5



To the *cis* isomer 26-4-1 (0.600 g, 1.33 mmol) in 4.4% HCOOH/CH<sub>3</sub>OH (50 ml) was added 10% Pd/C (0.600 g). The reaction mixture was stirred at R.T. under argon for 16 h, then filtered through celite and concentrated. The residue was purified by PTLC (1:10 (2M NH<sub>3</sub>/CH<sub>3</sub>OH)/CH<sub>2</sub>Cl<sub>2</sub>) to afford the product (0.230 g, 85%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.50 (4H, s), 7.06 (2H, m), 6.70 (1H, m), 6.40 (1H, s), 4.20 (1H, m), 3.30 (1H), 3.00 (3H, s), 1.50-2.30 (10H, m). MS m/e 360 (M+H).


Step 6

To a mixture of the product of Step 5 (0.140 g, 0.390 mmol) and 1M K<sub>2</sub>CO<sub>3</sub> (1.2 ml, 1.2 mmol) in THF (5 ml) was added MeSO<sub>2</sub>Cl (0.178 g, 1.55 mmol). The reaction mixture was stirred at R.T. for 16 h then subjected to PTLC (1:10 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (0.135 g, 79%). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz) δ 7.53 (4H, m), 7.20 (2H, m), 6.90 (1H, m), 4.10 (1H, m), 3.60 (1H, m), 2.90 (6H, s), 1.50-2.10 (8H, m). MS m/e 438 (M+H).

Example 27


A mixture of 26-3-1 (0.21 g, 0.59 mmol), hydroxylamine hydrochloride (0.82 g, 12 mmol), and sodium acetate (0.97 g, 12 mmol) in absolute EtOH (10 ml) was stirred at R.T. for 64 h. The mixture was partitioned between CH<sub>2</sub>Cl<sub>2</sub> (100 ml) and water (75 ml). The aqueous layer was extracted again with CH<sub>2</sub>Cl<sub>2</sub> (50 ml). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The residue was subjected to PTLC (1:19 CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product (210 mg, 95%). <sup>1</sup>H NMR (CD<sub>3</sub>OD, 400 MHz) δ 7.4-7.6 (4H, m), 7.20 (2H, m), 6.85 (1H, m), 4.39 (1H, m), 3.45 (1H, m), 2.90 (3H, s), 2.45 (1H, m), 2.28 (1H, m), 1.6-2.0 (5H, m). MS (ES) m/e 374 (M+H).

Use of the appropriate starting material and essentially the same procedure afforded the following compound.




MS (ES) m/e 388 (M+H).

51



## Step 1



To a mixture of 1-3-5 (100 mg, 0.31 mmol), 1 M NaOH (0.5 ml), and 1 M Na<sub>2</sub>CO<sub>3</sub> (0.5 ml) in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) was added 2-chloroethylsulfonate (100 mg, 0.61 mmol), and the reaction mixture was stirred for 16 hr. The reaction mixture was partitioned between water (25 ml) and CH<sub>2</sub>Cl<sub>2</sub> (25 ml). The organic layer was dried (MgSO<sub>4</sub>), filtered, and concentrated. Subjection of the residue to PTLC (1:4 acetone/CH<sub>2</sub>Cl<sub>2</sub>) gave the product (40 mg, 31%). MS (ES) m/e 418 (M+H).


10

## Step 2

To a stirred solution of the product of Step 1 (28-1-1) (50 mg, 0.12 mmol) in THF (10 ml) was added tetrabutylammonium hydroxide (0.5 g) in water (2 ml). After 16 hr, the reaction mixture was partitioned between water (25 ml) and CH<sub>2</sub>Cl<sub>2</sub> (100 ml). The organic layer was dried (MgSO<sub>4</sub>), filtered, and concentrated. Subjection of the residue to PTLC (5:95 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave the product (24 mg, 46%). HRMS calc. for C<sub>21</sub>H<sub>27</sub>FN<sub>3</sub>O<sub>4</sub>S (M+H) 436.1706. Found 436.1711.

15

## Example 29




20

To a solution of 1-3-1 (400 mg, 1.22 mmol) in DMF (5 ml) was added EDCI (25 mg, 1.30 mmol) and 1-cyano-3-methylisothiourea sodium salt (175 mg, 1.27 mmol). The reaction mixture was stirred for 16 h, then diluted with EtOAc (50 ml). The mixture was washed with water (10 ml), sat'd NaHCO<sub>3</sub> (20 ml) and water (10 ml). The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated. Subjection of the residue

25

to flash chromatography (gradient; 3:97 – 7:93 MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave the product (250 mg, 50%). HRMS calc. For C<sub>22</sub>H<sub>26</sub>N<sub>6</sub>OF (M+H) 409.2152. Found 409.2155.

Example 30



5

To a solution of 1-3-1 (500 mg, 1.53 mmol) in acetonitrile (10 ml) was added dimethyl-N-cyanodithioiminocarbonate (0.8 g, 5.5 mmol) and the reaction mixture was refluxed for 16 h. The reaction mixture was poured into water (50 ml) and extracted with EtOAc (50 ml). The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated.

10 Subjection of the residue to flash chromatography (1:2 acetone/hexanes) gave the product (150 mg, 24%). MS m/e 426.1 (M+H).

Method for Screening Compound 14 of  
Example 14 for Y5 Antagonist Activity In Vivo

15

Adult male Long-Evans or Sprague-Dawley rats (200-250 g, Charles River, MA) were maintained in individual cages at 22°C on a 12 hr light/12 hr dark cycle with lights on at 0400. Rats had free access to food (Teklad Lab Rodent Chow, Bartonville, IL) and water. All studies were conducted in an AAALAC accredited

20 facility following protocols approved by the Animal Care and Use Committee of the Schering-Plough Research Institute. The procedures were performed in accordance with the principles and guidelines established by the NIH for the care and use of laboratory animals.













Rats were anesthetized by intramuscular injection of a mixture of ketamine and












25 xylazine (100 and 10 mg/kg, respectively). A 22 gauge stainless steel cannula was stereotactically implanted into the lateral ventricle using the following coordinates: 1 mm posterior to bregma, 1.5 mm lateral to midline, 3.6 mm ventral to dura. After a three week recovery period, all animals were tested for correct cannula placement by intracerebroventricular (icv) infusion of human NPY (0.3 nmol). Only animals

30 demonstrating a profound feeding effect (>2 g) within 60 min of the infusion were retained for the study. Four groups of twelve animals were used in each study. Each group was balanced such that the average baseline and NPY-induced food intake













values were similar for each group. One group received an oral dose of vehicle while the other three groups received oral doses of the Y5 antagonist 14 one hour before icv administration of D-Trp34-NPY. D-Trp34-NPY was dissolved in 0.9% sterile saline (Sigma, St. Louis, MO) and were infused icv with a Hamilton infusion pump and  
5 syringe (Hamilton, Reno, NV) at a rate of 5  $\mu$ l/min. The guide cannula remained inserted for an additional minute to prevent diffusion up the needle track. The chow-filled feeder was weighed during the infusion period and then returned to the home cage with the animal immediately following treatment. Food consumption was monitored at 60, 120 and 240 min after icv infusion of peptides. Differences in food  
10 intake between groups were determined by analysis of variance followed by Dunnett's multiple comparison test. Compound 14 (0.1, 0.3, 1, and 3 mg/kg) dose responsively inhibited D-Trp34-NPY stimulated food intake with an ID50 of 0.5 mg/kg.













It will be recognized that the following examples can be prepared by adapting  
15 appropriate procedures described in Examples 1 – 30, or by applying methods known to those skilled in the art:








Example	Structure	MSm/e (M+H)
31		483
32		449
33		483
34		467
35		440
36		483
37		483
38		457
39		457
40		449
41		449
42		440













43		483
44		483
45		422
46		410
47		424
48		438
49		438
50		436
51		472
52		374
53		388



























54		402
55		402
56		400
57		442
58		414
59		428
60		396
61		403
62		431
63		414
64		423
65		360



66		374
67		388
68		388
69		410
70		424
71		422
72		424
73		386
74		404
75		356
76		370
77		392













78		406
79		420
80		418
81		420
82		384
83		384
84		382

85		388
86		466
87		531
88		452
89		467
90		452
91		428
92		402
93		416
94		416
95		430
96		456













97		456
98		430
99		442
100		480
101		444
102		467
103		465
104		465
105		428
106		465
107		422
108		410













109		424
110		438
111		438
112		436
113		472
114		374
115		400
116		388
117		402
118		402
119		442
120		414













121		428
122		408













123		431
124		338
125		352
126		428
127		396
128		368
129		395
130		435
131		437
132		407
133		443
134		449





135		381
136		450
137		388
138		402
139		416
140		417
141		450
142		464
143		416
144		389
145		442
146		356









147		370
148		403
149		371
150		389
151		449
152		385
153		449
154		449
155		511
156		449
157		519
158		465









159		467
160		501
161		511
162		466
163		467
164		466
165		449
166		449
167		447
168		531
169		448
170		448













171		452
172		466
173		467
174		468
175		440
176		452
177		450
178		422
179		434
180		434
181		448
182		449

68













183		403
184		487













185		459
186		487
187		409
188		420
189		436
190		401
191		435
192		485

193		449
194		523
195		463
196		450
197		442
198		420
199		438
200		427




201		387.1
202		388.1
203		387.1
204		386.1
205		393.1
206		323.1
207		465.1, 467.1
208		378.1
209		378.1
210		387.1
211		455.1
212		455.1























213		416.1
214		403.1
215		401.1
216		405.1
217		441.1
218		423.1
219		457.1
220		439.1
221		437.1
222		448.1
223		450.1
224		432.1











225		436.1
226		422.1
227		439.1
228		436.1
229		422.1
230		512.1
231		422.1
232		462.1
233		385
234		440.1
235		462.1
236		440.1











74











237		454.1
238		468.1
239		468.1

240		441
241		473
242		405
243		437
244		491
245		491
246		405
247		423
248		423
249		439











250		416
251		405
252		421
253		453
254		491
255		501
256		517
257		430
258		458
259		492













260		599
261		424
262		487
263		442
264		442
265		424
266		436
267		422
268		424
269		424













270		436
271		466
272		422
273		424
274		424
275		424
276		424
277		458
278		424
279		446













280		388
281		418
282		402
283		466
284		466
285		529
286		507
287		471
288		422
289		456









290		456
291		413
292		451
293		413
294		416
295		416
296		456
297		442
298		414
299		454

300		414
301		414
302		360
303		438
304		452
305		466
306		452
307		466
308		402
309		416
310		428
311		465


312		465
314		465
315		403
316		437
317		437
318		458
319		424
320		374
321		374
322		529
323		416
324		490

325		439
326		452
327		424
328		439
329		424
330		451
331		451
332		446
333		402
334		451
335		388
336		446

337		446
338		388
339		402
340		388
341		402
342		464


What is Claimed:

1. A compound having the structural formula I:





including its N-oxides, wherein

Y is



$R^1$  is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

$R^2$  is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>9</sub>)cycloalkyl or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl;



Z is

$OR^{10}$ ,  $-N(R^9)(R^{10})$  or  $-NH_2$ ;

j is 0, 1 or 2;

k is 1 or 2;

l is 0, 1 or 2;

m is 0, 1 or 2;

$R^4$  is 1-3 substituents independently selected from the group consisting of H, -OH, halogen, haloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -CN,  $-O(C_1-C_6)alkyl$ ,  $-O(C_3-C_7)cycloalkyl$ ,  $-O(C_1-C_6)alkyl(C_3-C_7)cycloalkyl$ ,  $-S(C_1-C_6)alkyl$ ,  $-S(C_3-C_7)cycloalkyl$ ,  $-S(C_1-C_6)alkyl(C_3-C_7)cycloalkyl$ ,  $-NH_2$ ,  $-NR^9R^{10}$ ,  $-NO_2$ ,  $-CONH_2$ ,  $-CONR^9R^{10}$  and  $NR^2COR^{10}$ ;

$R^5$  is 1-3 substituents independently selected from the group consisting of H, halogen, -OH, haloalkyl, haloalkoxy, -CN,  $-NO_2$ , (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,

(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
-O(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -CONH<sub>2</sub> and -CONR<sup>9</sup>R<sup>10</sup>;

- R<sup>6</sup> is -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO<sub>2</sub>(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
-SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)haloalkyl, -SO<sub>2</sub>(hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkyl),  
5 -SO<sub>2</sub>(amino(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>(alkoxy(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>(alkylamino(C<sub>2</sub>-C<sub>6</sub>)alkyl),  
-SO<sub>2</sub>(dialkylamino(C<sub>2</sub>-C<sub>6</sub>)alkyl), -SO<sub>2</sub>(aryl), -SO<sub>2</sub>(heteroaryl), -SO<sub>2</sub>(aryl(C<sub>2</sub>-C<sub>6</sub>)alkyl),  
-SO<sub>2</sub>NH<sub>2</sub>, -SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup>, -C(O)(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
-C(O)(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)aryl, -C(O)heteroaryl, -C(O)NR<sup>9</sup>R<sup>10</sup>,  
-C(O)NH<sub>2</sub>, -C(S)NR<sup>9</sup>R<sup>10</sup>, -C(S)NH<sub>2</sub>, aryl, heteroaryl, -(CH<sub>2</sub>)<sub>n</sub>C(O)NH<sub>2</sub>,  
10 - (CH<sub>2</sub>)<sub>n</sub>C(O)NR<sup>9</sup>R<sup>10</sup>, -C(=NCN)alkylthio, -C(=NCN)NR<sup>9</sup>R<sup>10</sup>, (C<sub>1</sub>-C<sub>6</sub>)alkyl,  
(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl  
or -C(O)OR<sup>9</sup>, n = 1 to 6;

R<sup>7</sup> = H or alkyl;

- R<sup>8</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl;  
15 heteroaryl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO<sub>2</sub>(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl,  
-SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)haloalkyl or -SO<sub>2</sub>(aryl);

R<sup>9</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or heteroaryl; and,


- R<sup>10</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
20 aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or heteroaryl;

or R<sup>9</sup> and R<sup>10</sup> taken together can form a 4-7 membered ring containing 1 or 2  
heteroatoms;

- or its pharmaceutically acceptable addition salt and/or hydrate thereof, or  
prodrug thereof, or where applicable, a geometric or optical isomer or a racemic  
25 mixture thereof.

87


2. A compound of claim 1 wherein





5 3. A compound of claim 2 wherein R<sup>5</sup> is 1-3 substituents independently selected from the group consisting of H, halogen, haloalkyl, alkoxy and haloalkoxy and the sum of j and k is 1, 2 or 3.

4. A compound of claim 2 wherein R<sup>6</sup> is SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, SO<sub>2</sub>hydroxy(C<sub>2</sub>-C<sub>6</sub>)alkyl,  
10 SO<sub>2</sub>(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, SO<sub>2</sub>NR<sup>9</sup>R<sup>10</sup> or SO<sub>2</sub>NH<sub>2</sub>.

5. A compound of claim 1 selected from the group consisting of





15





20







5




10



and their

pharmaceutically acceptable addition salts and/or hydrates thereof, or prodrugs thereof, or where applicable, geometric or optical isomers or a racemic mixtures thereof.

6. A compound of claim 1, wherein the compound is





5



or its pharmaceutically acceptable addition salt and/or hydrate thereof, or prodrug thereof, or where applicable, a geometric or optical isomer or a racemic mixture thereof.

- 10 7. A compound of claim 2 wherein  $R^6$  is  $C(O)$ heteroraryl,  $C(O)(C_1-C_6)$  alkyl or  $C(O)(C_3-C_7)$ cycloalkyl.

8. A compound of claim 1 selected from the group consisting of







15





20

90





5




and their pharmaceutically acceptable addition salts and/or hydrates thereof, or  
 10 prodrugs thereof, or where applicable, geometric or optical isomers or a racemic  
 mixtures thereof.

9. A compound of claim 2 wherein R<sup>6</sup> is heteroaryl.

15 10. A compound of claim 1 selected from the group consisting of




91




- and their pharmaceutically acceptable addition salts and/or hydrates thereof, or prodrugs thereof, or where applicable, geometric or optical isomers or a racemic mixtures thereof.

11. A compound of claim 1 wherein



10

12. A compound of claim 11 wherein  $R^5$  is 1 to 3 substituents independently selected from the group consisting of H, halogen, haloalkyl and haloalkoxy and the sum of j and k is 1, 2 or 3.
13. A compound of claim 11 wherein  $R^6$  is  $\text{SO}_2(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $\text{SO}_2(\text{C}_3\text{-C}_7)\text{cycloalkyl}$ ,  $\text{SO}_2\text{NR}^9\text{R}^{10}$  or  $\text{SO}_2\text{NH}_2$ .
14. A compound of the formula




20

or its pharmaceutically acceptable addition salt and/or hydrate thereof, or prodrug thereof, or where applicable, a geometric or optical isomer or a racemic mixture thereof.

15. A compound of claim 11 wherein  $R^6$  is  $C(O)heteroaryl$ ,  $C(O)(C_1-C_6) alkyl$  or  $C(O)(C_3-C_7)cycloalkyl$ .

16. A compound of claim 1 selected from the group consisting of

5



and their pharmaceutically acceptable addition salts and/or hydrates thereof, or prodrugs thereof, or where applicable, geometric or optical isomers or a racemic mixtures thereof.

























10





















17. A compound of claim 11 wherein  $R^8$  is heteroaryl.





















18. A compound of claim 1 selected from the group consisting of those having the structural formulas set forth in the following table, and the pharmaceutically

15 acceptable addition salts and/or hydrates thereof, or prodrugs thereof, or where applicable, geometric or optical isomers or a racemic mixtures thereof:























Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H






















Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H



















Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H

Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		2-F
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H



Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H

Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>	-CH <sub>2</sub> CONH <sub>2</sub>	-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H

Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H
	-H	-CH <sub>3</sub>		-H

19. The compound of claim 1 selected from the compounds of Examples: 29-59, 61-90, 95-216, 218-219, 221-262, 265, 267, 269-294, 296-297, 299-326, 328-337, 340-342 and their pharmaceutically acceptable addition salts and/or hydrates thereof,

or prodrugs thereof, or where applicable, geometric or optical isomers or a racemic mixtures thereof.

20. A pharmaceutical composition comprising a compound of formula I as defined  
5 in claim 1 in combination with a pharmaceutically acceptable carrier.

21. A method of treating obesity, an eating disorder or diabetes comprising  
administering an effective amount of a compound of formula 1 as defined in Claim 1 to  
a mammal in need of such treatment. A pharmaceutical composition, which comprises  
10 an effective amount of a compound as, defined in claim 1 and a pharmaceutically  
acceptable carrier thereof.

22. A method of treating metabolic or eating disorders comprising administering to  
a mammal in need of such treatment a therapeutically effective amount of a  
15 compound of claim 1 or a prodrug thereof or a pharmaceutically acceptable salt of  
said compound or of said prodrug.

23. The method of claim 22 wherein said metabolic disorder is obesity.

24. The method of claim 22 wherein said eating disorder is hyperphagia.

20 25. A method of treating disorders associated with obesity comprising  
administering to a mammal in need of such treatment a therapeutically effective  
amount of a compound of claim 1 or a prodrug thereof or a pharmaceutically  
acceptable salt of said compound or of said prodrug.

25 26. The method of claim 25 wherein said disorders associated with obesity are  
Type II Diabetes, insulin resistance, hyperlipidemia and hypertension.

27. A pharmaceutical composition which comprises a therapeutically effective  
amount of a composition comprising:

a first compound, said first compound being a compound of claim 1, a prodrug  
30 thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug;

a second compound, said second compound being an anti-obesity and/or anorectic agent such as a  $\beta_3$  agonist, a thryomimetic agent, an anorectic agent or an NPY antagonist; and

a pharmaceutically acceptable carrier thereof.

5

28. A method of treating a metabolic or eating disorder which comprises administering to a mammal in need of such treatment

an amount of a first compound, said first compound being a compound of claim 1, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug;

10

a second compound, said second compound being an antiobesity and/or anorectic agent such as a  $\beta_3$  agonist, a thryomimetic agent, an anorectic agent or an NPY antagonist;

wherein the amounts of the first and second compounds result in a therapeutic effect.

15

29. A pharmaceutical composition which comprises a therapeutically effective amount of a composition comprising:

a first compound, said first compound being a compound of claim 1, a prodrug thereof, or a pharmaceutically acceptable salt of said compound or of said prodrug;

20

a second compound, said second compound being an aldose reductase inhibitor, a glycogen phosphorylase inhibitor, a sorbitol dehydrogenase inhibitor, a protein tyrosine phosphatase 1B inhibitor, a dipeptidyl protease inhibitor, insulin (including orally bioavailable insulin preparations), an insulin mimetic, metformin, acarbose, a PPAR-gamma ligand such as troglitazone, rosiglitazone, pioglitazone, or GW-1929, a sulfonylurea, glipazide, glyburide, or chlorpropamide; and

25

a pharmaceutically acceptable carrier therefor.

30. A pharmaceutical composition made by combining the compound of claim 1 and a pharmaceutically acceptable carrier therefor.

31. A process for making a pharmaceutical composition comprising combining a compound of claim 1 and a pharmaceutically acceptable carrier.

30

**BLANK PAGE**

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number  
WO 02/22592 A3(51) International Patent Classification<sup>7</sup>: C07D 211/58,  
A61K 31/4465, 31/4525, 31/4535, 31/4545, A61P 3/04,  
C07D 211/96, 401/04, 401/06, 409/12, 409/14, 417/04,  
C07C 275/30, 311/07

(21) International Application Number: PCT/US01/28324

(22) International Filing Date:  
12 September 2001 (12.09.2001)








(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/232,255 14 September 2000 (14.09.2000) US(71) Applicant: SCHERING CORPORATION [US/US]:  
Patent Department - K-6-1 1990, 2000 Galloping Hill  
Road, Kenilworth, NJ 07033-0530 (US).(72) Inventors: GREENLEE, William, J.; 115 Herrick  
Avenue, Teaneck, NJ 07666 (US). HUANG, Ying; 8Ivy Terrace, East Brunswick, NJ 08816 (US). KELLY,  
Joseph, M.; 112 Princeton Road, Parlin, NJ 08859 (US).  
MC COMBIE, Stuart, W.; 28 Hanford Place, Caldwell,  
NJ 07006 (US). STAMFORD, Andrew, W.; 27 Overlook  
Road, Chatham Township, NJ 07928 (US). WU, Yusheng;  
235 East 40th Street, Apt. #20C, New York, NY 10016  
(US).(74) Agent: LEE, William, Y.; Schering-Plough Corporation,  
Patent Department - K-6-1 1990, 2000 Galloping Hill Road,  
Kenilworth, NJ 07033-0530 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,  
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,  
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT,  
RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ,  
VN, YU, ZA.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

[Continued on next page]

(54) Title: SUBSTITUTED UREA NEUROPEPTIDE Y Y5 RECEPTOR ANTAGONISTS



(57) Abstract: Compounds represented by structural formula (I) including its N-oxides wherein Y is (a) or (b); R<sup>1</sup> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl; R<sup>2</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>9</sub>)cycloalkyl or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl; R<sup>3</sup> is (c), (d), (e), (f), (g), (h) or (i); R<sup>4</sup> is 1-3 substituents independently selected from the group consisting of H, -OH, halogen, haloalkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, -CN, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -S(C<sub>1</sub>-C<sub>6</sub>)alkyl, -S(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -S(C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, -NH<sub>2</sub>, -NR<sup>9</sup>R<sup>10</sup>, -NO<sub>2</sub>, -CONH<sub>2</sub>, -CONR<sup>9</sup>R<sup>10</sup> and NR<sup>9</sup>COR<sup>10</sup>; or where applicable, a geometric or optical isomer or a racemic mixture thereof, are claimed, as well as additional novel compounds; also claimed are pharmaceutical compositions and methods of using the aforesaid compounds in the treatment of obesity, eating disorders such as hyperphagia and diabetes.



IT, LU, MC, NL, PT, SE, TR). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) Date of publication of the international search report:  
27 June 2002

**Published:**

- with international search report  
before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



# INTERNATIONAL SEARCH REPORT

Int. Application No

PCT/US 01/28324

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D211/58 A61K31/4465 A61K31/4525 A61K31/4535 A61K31/4545  
A61P3/04 C07D211/96 C07D401/04 C07D401/06 C07D409/12  
C07D409/14 C07D417/04 C07C275/30 C07C311/07

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99 64394 A (STAMFORD ANDREW W ;DUGAR SUNDEEP (US); SCHERING CORP (US); WU YUSH) 16 December 1999 (1999-12-16) page 53 -page 63; claims page 8, line 16 - line 18 page 47 -page 48; examples 19E,19F,19K ---	1-31
Y	WO 97 19682 A (FINN JOHN M ;JEON YOON T (US); PING DU (US); DHANOA DALJIT S (US);) 5 June 1997 (1997-06-05) page 131 -page 138; claim 1 page 6, line 2 - line 12 page 76 -page 77; example 87 ---	1-31
A	WO 98 35957 A (BAYER AG) 20 August 1998 (1998-08-20) page 53 -page 55; claims 1-7 page 38; table 2, compound no. 271 ---	1-31
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*8\* document member of the same patent family

Date of the actual completion of the international search

21 March 2002

Date of mailing of the international search report

18/04/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fink, D

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/28324

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MCNALLY J J ET AL: "N-Acylated alpha-(3-pyridylmethyl)-beta-aminotetralin antagonists of the human neuropeptide Y Y5 receptor" BIOORGANIC &amp; MEDICINAL CHEMISTRY LETTERS, vol. 10, no. 15, 7 August 2000 (2000-08-07), pages 1641-1643, XP004213213 ISSN: 0960-894X the whole document -----</p>	1-31

# INTERNATIONAL SEARCH REPORT

Information on patent family members

In tional Application No

PCT/US 01/28324

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9964394	A	16-12-1999	AU 4317899 A CN 1311773 T EP 1086078 A1 WO 9964394 A1	30-12-1999 05-09-2001 28-03-2001 16-12-1999
WO 9719682	A	05-06-1997	AU 1328197 A US 6211241 B1 WO 9719682 A1	19-06-1997 03-04-2001 05-06-1997
WO 9835957	A	20-08-1998	AU 6144098 A CA 2251368 A1 EP 0910565 A1 JP 2000510164 T US 5939462 A WO 9835957 A1	08-09-1998 20-08-1998 28-04-1999 08-08-2000 17-08-1999 20-08-1998

**BLANK PAGE**